




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<div>(51) International Patent Classification⁶ : C12N 15/53, 15/83, 5/10, C12P 7/64, C11B 1/00, A61K 31/20, A23L 1/30, A23K 1/00</div>	<div>A1</div>	<div>(11) International Publication Number: WO 98/46765</div> <div>(43) International Publication Date: 22 October 1998 (22.10.98)</div>
<div>(21) International Application Number: PCT/US98/07422</div> <div>(22) International Filing Date: 10 April 1998 (10.04.98)</div> <div>(30) Priority Data: 08/833,610 11 April 1997 (11.04.97) US</div> <div>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/833,610 (CIP) Filed on 11 April 1997 (11.04.97)</div> <div>(71) Applicants (for all designated States except US): CALGENE LLC [US/US]; 1920 Fifth Street, Davis, CA 95616 (US). ABOIT LABORATORIES [US/US]; 100 Abbott Park Road, Abbott Park, IL 60064-3500 (US).</div> <div>(72) Inventors; and</div> <div>(75) Inventors/Applicants (for US only): KNUTZON, Deborah [US/US]; 6110 Rockhurst Way, Granite Bay, CA 95746 (US). MUKERJI, Pradip [US/US]; 1069 Arcaro Drive, Gahanna, OH 43230 (US). HUANG, Yung-Sheng [CA/US]; 2462 Danvers Court, Upper Arlington, OH 43220 (US). THURMOND, Jennifer [US/US]; 3702 Adirondack, Colum-</div>	<div>bus, OH 43231 (US). CHAUDHARY, Sunita [IN/US]; 3419 Woodbine Place, Pearland, TX 77584 (US). LEONARD, Amanda, Eun-Yeong [US/US]; 581 Shadewood Court, Gahanna, OH 43230 (US).</div> <div>(74) Agents: WARD, Michael, R. et al.; Limbach & Limbach L.L.P., 2001 Ferry Building, San Francisco, CA 94111-4262 (US).</div> <div>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</div> <div>Published</div> <div>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</div>	

(54) Title: METHODS AND COMPOSITIONS FOR SYNTHESIS OF LONG CHAIN POLYUNSATURATED FATTY ACIDS

(57) Abstract

The present invention relates to a fatty acid $\Delta 5$ -desaturase able to catalyze the conversion of dihomo- γ -linolenic acid to arachidonic acid. Nucleic acid sequences encoding $\Delta 5$ -desaturase, nucleic acid sequences which hybridize thereto, DNA constructs comprising a $\Delta 5$ -desaturase gene, and recombinant host microorganism or animal expressing increased levels of a $\Delta 5$ -desaturase are described. Methods for desaturating a fatty acid at the $\Delta 5$ position and for producing arachidonic acid by expressing increased levels of a $\Delta 5$ -desaturase are disclosed. Fatty acids, and oils containing them, which have been desaturated by a $\Delta 5$ -desaturase produced by recombinant host microorganisms or animals are provided. Pharmaceutical compositions, infant formulas or dietary supplements containing fatty acids which have been desaturated by a $\Delta 5$ -desaturase produced by a recombinant host microorganism or animal also are described.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS AND COMPOSITIONS FOR SYNTHESIS OF LONG CHAIN POLYUNSATURATED FATTY ACIDS

RELATED APPLICATION

5 This application is a continuation in part application of Serial Number 08/833,610 filed April 11, 1997.

INTRODUCTION

Field of the Invention

 This invention relates to modulating levels of enzymes and/or enzyme
10 components relating to production of long chain poly-unsaturated fatty acids (PUFAs) in a microorganism or animal.

Background

 Two main families of polyunsaturated fatty acids (PUFAs) are the $\omega 3$
15 fatty acids, exemplified by eicosapentaenoic acid (EPA), and the $\omega 6$ fatty acids, exemplified by arachidonic acid (ARA). PUFAs are important components of the plasma membrane of the cell, where they may be found in such forms as phospholipids. PUFAs are necessary for proper development, particularly in the developing infant brain, and for tissue formation and repair. PUFAs also serve
20 as precursors to other molecules of importance in human beings and animals, including the prostacyclins, eicosanoids, leukotrienes and prostaglandins.

 Four major long chain PUFAs of importance include docosahexaenoic acid (DHA) and EPA, which are primarily found in different types of fish oil, gamma-linolenic acid (GLA), which is found in the seeds of a number of plants,
25 including evening primrose (*Oenothera biennis*), borage (*Borago officinalis*) and black currants (*Ribes nigrum*), and stearidonic acid (SDA), which is found in marine oils and plant seeds. Both GLA and another important long chain PUFA, arachidonic acid (ARA), are found in filamentous fungi. ARA can be

purified from animal tissues including liver and adrenal gland. GLA, ARA, EPA and SDA are themselves, or are dietary precursors to, important long chain fatty acids involved in prostaglandin synthesis, in treatment of heart disease, and in development of brain tissue.

5 Polyunsaturated fatty acids have a number of pharmaceutical and medical applications including treatment of heart disease, cancer and arthritis.

For DHA, a number of sources exist for commercial production including a variety of marine organisms, oils obtained from cold water marine fish, and egg yolk fractions. For ARA, microorganisms including the genera
10 *Mortierella*, *Entomophthora*, *Phytium* and *Porphyridium* can be used for commercial production. Commercial sources of SDA include the genera *Trichodesma* and *Echium*. Commercial sources of GLA include evening primrose, black currants and borage. However, there are several disadvantages associated with commercial production of PUFAs from natural sources. Natural
15 sources of PUFAs, such as animals and plants, tend to have highly heterogeneous oil compositions. The oils obtained from these sources therefore can require extensive purification to separate out one or more desired PUFAs or to produce an oil which is enriched in one or more PUFA. Natural sources also are subject to uncontrollable fluctuations in availability. Fish stocks may
20 undergo natural variation or may be depleted by overfishing. Fish oils have unpleasant tastes and odors, which may be impossible to economically separate from the desired product, and can render such products unacceptable as food supplements. Animal oils, and particularly fish oils, can accumulate environmental pollutants. Weather and disease can cause fluctuation in yields
25 from both fish and plant sources. Cropland available for production of alternate oil-producing crops is subject to competition from the steady expansion of human populations and the associated increased need for food production on the remaining arable land. Crops which do produce PUFAs, such as borage, have not been adapted to commercial growth and may not perform well in
30 monoculture. Growth of such crops is thus not economically competitive where more profitable and better established crops can be grown. Large scale

fermentation of organisms such as *Mortierella* is also expensive. Natural animal tissues contain low amounts of ARA and are difficult to process. Microorganisms such as *Porphyridium* and *Mortierella* are difficult to cultivate on a commercial scale.

5 Dietary supplements and pharmaceutical formulations containing PUFAs can retain the disadvantages of the PUFA source. Supplements such as fish oil capsules can contain low levels of the particular desired component and thus require large dosages. High dosages result in ingestion of high levels of undesired components, including contaminants. Unpleasant tastes and odors of
10 the supplements can make such regimens undesirable, and may inhibit compliance by the patient. Care must be taken in providing fatty acid supplements, as overaddition may result in suppression of endogenous biosynthetic pathways and lead to competition with other necessary fatty acids in various lipid fractions *in vivo*, leading to undesirable results. For example,
15 Eskimos having a diet high in ω 3 fatty acids have an increased tendency to bleed (U.S. Pat. No. 4,874,603).

A number of enzymes are involved in PUFA biosynthesis. Linolenic acid (LA, 18:2 Δ 9, 12) is produced from oleic acid (18:1 Δ^0) by a Δ 12-desaturase. GLA (18:3 Δ 6, 9, 12) is produced from linoleic acid (LA, 18:2 Δ 9,
20 12) by a Δ 6-desaturase. ARA (20:4 Δ 5, 8, 11, 14) production from dihomo-gamma-linolenic acid (DGLA, 20:3 Δ 8, 11, 14) is catalyzed by a Δ 5-desaturase. However, animals cannot desaturate beyond the Δ 9 position and therefore cannot convert oleic acid (18:1 Δ 9) into linolenic acid (18:2 Δ 9, 12). Likewise, α -linoleic acid (ALA, 18:3 Δ 9, 12, 15) cannot be synthesized by mammals.
25 Other eukaryotes, including fungi and plants, have enzymes which desaturate at positions Δ 12 and Δ 15. The major poly-unsaturated fatty acids of animals therefore are either derived from diet and/or from desaturation and elongation of linoleic acid (18:2 Δ 9, 12) or α -linolenic acid (18:3 Δ 9, 12, 15). Therefore it is of interest to obtain genetic material involved in PUFA biosynthesis from
30 species that naturally produce these fatty acids and to express the isolated

material in a microbial or animal system which can be manipulated to provide production of commercial quantities of one or more PUFAs. Thus there is a need for fatty acid desaturases, genes encoding them, and recombinant methods of producing them. A need further exists for oils containing higher relative proportions of and/or enriched in specific PUFAs. A need also exists for reliable economical methods of producing specific PUFAs.

Relevant Literature

Production of gamma-linolenic acid by a $\Delta 6$ -desaturase is described in USPN 5,552,306. Production of 8, 11-eicosadienoic acid using *Mortierella alpina* is disclosed in USPN 5,376,541. Production of docosahexaenoic acid by dinoflagellates is described in USPN 5,407,957. Cloning of a $\Delta 6$ -palmitoyl-acyl carrier protein desaturase is described in PCT publication WO 96/13591 and USPN 5,614,400. Cloning of a $\Delta 6$ -desaturase from borage is described in PCT publication WO 96/21022. Cloning of $\Delta 9$ -desaturases is described in the published patent applications PCT WO 91/13972, EP 0 550 162 A1, EP 0 561 569 A2, EP 0 644 263 A2, and EP 0 736 598 A1, and in USPN 5,057,419. Cloning of $\Delta 12$ -desaturases from various organisms is described in PCT publication WO 94/11516 and USPN 5,443,974. Cloning of $\Delta 15$ -desaturases from various organisms is described in PCT publication WO 93/11245. All publications and U.S. patents or applications referred to herein are hereby incorporated in their entirety by reference.

Summary of the Invention

Novel compositions and methods are provided for preparation of polyunsaturated long chain fatty acids or PUFAs. The compositions include nucleic acids encoding a $\Delta 5$ -desaturase and/or polypeptides having $\Delta 5$ -desaturase activity, the polypeptides, and probes for isolating and detecting the same. The methods involve growing a host microorganism or animal which contains and expresses one or more transgenes encoding a $\Delta 5$ -desaturase and/or a polypeptide having $\Delta 5$ -desaturase activity. Expression of the desaturase

polypeptide provides for a relative increase in $\Delta 5$ -desaturated PUFA, or metabolic progeny therefrom, as a result of altered concentrations of enzymes and substrates involved in PUFA biosynthesis. The invention finds use for example in the large scale production of PUFA containing oils which include,
5 for example, ARA, EPA and/or DHA.

In a preferred embodiment, a nucleic acid sequence comprising a $\Delta 5$ -desaturase depicted in Figure 3A-D (SEQ ID NO 1), a polypeptide encoded by the nucleic acid, and a purified or isolated polypeptide depicted in Figure 3A-D (SEQ ID NO: 2), and an isolated nucleic acid encoding the polypeptide of
10 Figure 3A-D (SEQ ID NO: 2) are provided. Another embodiment of the invention is an isolated nucleic acid sequence which encodes a polypeptide, wherein said polypeptide desaturates a fatty acid molecule at carbon 5 from the carboxyl end of the molecule. The nucleic acid is preferably derived from a eukaryotic cell, such as a fungal cell, or a fungal cell of the genus *Mortierella*,
15 or of the genus/species *Mortierella alpina*. Also preferred is an isolated nucleic acid comprising a sequence which anneals to a nucleotide sequence depicted in Figure 3A-3D (SEQ ID NO: 1), and a nucleic acid which encodes an amino acid sequence depicted in Figure 3A-D (SEQ ID NO: 2). In particular, the nucleic acid encodes an amino acid sequence depicted in Figure 3A-D (SEQ ID NO: 2)
20 which is selected from the group consisting of amino acid residues 30-38, 41-44, 171-175, 203-212, and 387-394. In an additional embodiment, the invention provides an isolated or purified polypeptide which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of the molecule. Also provided is an isolated nucleic acid sequence which hybridizes to a nucleotide sequence
25 depicted in Figure 3A-D (SEQ ID NO 1), an isolated nucleic acid sequence having at least about 50% identity to Figure 3A-D (SEQ ID NO 1).

The present invention further includes a nucleic acid construct comprising a nucleotide sequence depicted in a Figure 3A-D (SEQ ID NO: 1) linked to a heterologous nucleic acid; a nucleic acid construct comprising a
30 nucleotide sequence depicted in a Figure 3A-D (SEQ ID NO: 1) operably linked to a promoter; and a nucleic acid construct comprising a nucleotide sequence

depicted in a Figure 3A-D (SEQ ID NO: 1) operably linked to a promoter which is functional in a microbial cell. In a preferred embodiment, the microbial cell is a yeast cell, and the nucleotide sequence is derived from a fungus, such as a fungus of the genus *Mortierella*, particularly a fungus of the species *Mortierella alpina*.

In another embodiment of the invention, a nucleic acid construct is provided which comprises a nucleotide sequence which encodes a polypeptide comprising an amino acid sequence which corresponds to or is complementary to an amino acid sequence depicted in Figure 3A-D (SEQ ID NO: 2), wherein the nucleotide sequence is operably linked to a promoter which is functional in a host cell, and wherein the nucleotide sequence encodes a polypeptide which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of a fatty acid molecule. Additionally, provided by the invention is a nucleic acid construct comprising a nucleotide sequence which encodes a functionally active $\Delta 5$ -desaturase, where the desaturase includes an amino acid sequence which corresponds to or is complementary to all of or a portion of an amino acid sequence depicted in a Figure 3A-D (SEQ ID NO: 2), wherein the nucleotide sequence is operably linked to a promoter functional in a host cell.

The invention also includes a host cell comprising a nucleic acid construct of the invention. In a preferred embodiment, a recombinant host cell is provided which comprises at least one copy of a DNA sequence which encodes a functionally active *Mortierella alpina* fatty acid desaturase having an amino acid sequence as depicted in Figure 3A-D (SEQ ID NO: 2), wherein the cell or an ancestor of the cell was transformed with a vector comprising said DNA sequence, and wherein the DNA sequence is operably linked to a promoter. The host cell is either eukaryotic or prokaryotic. Preferred eukaryotic host cells are those selected from the group consisting of a mammalian cell, an insect cell, a fungal cell, and an algae cell. Preferred mammalian cells include an avian cell, a fungal cell such as a yeast, and a marine algae cell. Preferred prokaryotic cells include those selected from the group consisting of a bacteria, a cyanobacteria, cells which contain a

bacteriophage, and/or a virus. The DNA sequence of the recombinant host cell preferably contains a promoter which is functional in the host cell.

The host cells of the invention which contain the DNA sequences of the invention are enriched for fatty acids, such as 20:3 fatty acids. In a preferred
5 embodiment, the host cells are enriched for 20:4 fatty acids as compared to an untransformed host cell which is devoid of said DNA sequence, and/or enriched for 20:5 fatty acids compared to an untransformed host cell which is devoid of said DNA sequence. In yet another preferred embodiment, the invention provides a recombinant host cell which comprises a fatty acid selected from the
10 group consisting of a dihomo- γ -linolenic acid, n-6 eicosatrienoic acid, 20:3n-6 acid and 20:3 (8,11,14) acid.

The present invention also includes method for production of arachidonic acid in a microbial cell culture, where the method comprises growing a microbial cell culture having a plurality of microbial cells which
15 contain one or more nucleic acids encoding a polypeptide which converts dihomo- γ -linolenic acid to arachidonic acid, wherein the nucleic acid is operably linked to a promoter, under conditions whereby said one or more nucleic acids are expressed, whereby arachidonic acid is produced in the microbial cell culture. In several preferred embodiments of the invention, the
20 polypeptide is an enzyme which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of the fatty acid molecule; the nucleic acid is derived from a *Mortierella sp.*; and the substrate for said polypeptide is exogenously supplied. The microbial cells used in the methods can be either eukaryotic cells or prokaryotic cells. The preferred eukaryotic cells are those selected
25 from the group consisting of a mammalian cell, an insect cell, a fungal cell, and an algae cell. Preferred mammalian cells include an avian cell, a preferred fungal cell is a yeast, and the preferred algae cell is a marine algae cell. The preferred prokaryotic cells include those selected from the group consisting of a bacteria, a cyanobacteria, cells which contain a bacteriophage, and/or a virus.
30 The nucleic acid sequence encoding the polypeptide of the microbial cell preferably contains a promoter which is functional in the host cell which

optionally is an inducible promoter for example by components of the culture broth. The preferred microbial cells used in the methods are yeast cells, such as *Saccharomyces* cells.

5 In another embodiment of the invention, a recombinant yeast cell is provided which converts greater than about 5% of 20:3 fatty acid substrate to a 20:4 fatty acid product.

Also provided is an oil comprising one or more PUFA. The amount of said one or more PUFAs is approximately 0.3-30% arachidonic acid (ARA), approximately 0.2-30% dihomo- γ -linolenic acid (DGLA), and approximately
10 0.2-30% γ -linolenic acid (GLA). A preferred oil of the invention is one in which the ratio of ARA:DGLA:GLA is approximately 1.0:19.0:30 to 6.0:1.0:0.2. Another preferred embodiment of the invention is a pharmaceutical composition comprising the oils in a pharmaceutically acceptable carrier. Further provided is a nutritional composition comprising the oils of the
15 invention. The nutritional compositions of the invention preferably are administered to a mammalian host parenterally or internally. A preferred composition of the invention for internal consumption is an infant formula. In a preferred embodiment, the nutritional compositions of the invention are in a liquid form or a solid form.

20 The present invention also includes a method for desaturating a fatty acid, where the method comprises culturing a recombinant microbial cell of the invention under conditions suitable for expression of a polypeptide encoded by the nucleic acid, wherein the host cell further comprises a fatty acid substrate of the polypeptide. In a preferred embodiment, a fatty acid desaturated by the
25 methods is provided, including an oil comprising the fatty acid.

The present invention is also directed to purified nucleotide and peptide sequences presented in SEQ ID NO:1-34. The present invention is further directed toward methods of using the sequences presented in SEQ ID NO:1-34 as probes to identify related sequences, as components of expression systems
30 and as components of systems useful for producing transgenic oil.

The present invention is further directed to methods of obtaining altered long chain poly unsaturated fatty acid biosystems by growing transgenic microbes which encode transgene expression products which desaturate a fatty acid molecule at carbon 5 from the carboxyl end of the fatty acid molecule.

5 The present invention is further directed to formulas, dietary supplements or dietary supplements in the form of a liquid or a solid containing the long chain fatty acids of the invention. These formulas and supplements may be administered to a human or an animal.

10 The formulas and supplements of the invention may further comprise at least one macronutrient selected from the group consisting of coconut oil, soy oil, canola oil, mono- and diglycerides, glucose, edible lactose, electrodialysed whey, electrodialysed skim milk, milk whey, soy protein, and other protein hydrolysates.

15 The formulas of the present invention may further include at least one vitamin selected from the group consisting of Vitamins A, C, D, E, and B complex; and at least one mineral selected from the group consisting of calcium, magnesium, zinc, manganese, sodium, potassium, phosphorus, copper, chloride, iodine, selenium, and iron.

20 The present invention is further directed to a method of treating a patient having a condition caused by insufficient intake or production of polyunsaturated fatty acids comprising administering to the patient a dietary substitute of the invention in an amount sufficient to effect treatment of the patient.

The present invention is further directed to cosmetic and pharmaceutical compositions of the material of the invention.

25 The present invention is also directed to an isolated nucleotide sequence comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO:13; SEQ ID NO:15; SEQ ID NO:17; SEQ ID NO:19; SEQ ID NO:21; SEQ ID NO:22; SEQ ID NO:22; SEQ ID NO:24; SEQ ID NO:25; SEQ ID NO:26 and SEQ ID NO:27.

The present invention is also directed to an isolated peptide sequence comprising a peptide sequence selected from the group consisting of: SEQ ID NO:14; SEQ ID NO:16; SEQ ID NO:18; SEQ ID NO:20; SEQ ID NO:28; SEQ ID NO:29; SEQ ID NO:30; SEQ ID NO:31; SEQ ID NO:32; SEQ ID
5 NO:33 and SEQ ID NO:34.

The present invention is further directed to transgenic oils in pharmaceutically acceptable carriers. The present invention is further directed to nutritional supplements, cosmetic agents and infant formulae containing transgenic oils.

10 The present invention is further directed to a method for obtaining altered long chain polyunsaturated fatty acid biosynthesis comprising the steps of: growing a microbe having cells which contain a transgene which encodes a transgene expression product which desaturates a fatty acid molecule at carbon
5 from the carboxyl end of said fatty acid molecule, wherein the transgene is
15 operably associated with an expression control sequence, under conditions whereby the transgene is expressed, whereby long chain polyunsaturated fatty acid biosynthesis in the cells is altered.

The present invention is further directed to the use of chain polyunsaturated fatty acid selected from the group consisting of ARA, DGLA
20 and EPA.

The present invention is further directed toward pharmaceutical compositions comprising at least one nutrient selected from the group consisting of a vitamin, a mineral, a carbohydrate, a sugar, an amino acid, a free fatty acid, a phospholipid, an antioxidant, and a phenolic compound.

Brief Description of the Drawings

Figure 1 shows possible pathways for the synthesis of arachidonic acid (20:4 Δ 5, 8, 11, 14) and stearidonic acid (18:4 Δ 6, 9, 12, 15) from palmitic acid (C_{16}) from a variety of organisms, including algae, *Mortierella* and humans.

- 5 These PUFAs can serve as precursors to other molecules important for humans and other animals, including prostacyclins, leukotrienes, and prostaglandins, some of which are shown.

Figure 2 shows possible pathways for production of PUFAs in addition to ARA, including EPA and DHA, for a variety of organisms.

- 10 Figure 3A-D shows the DNA sequence of the *Mortierella alpina* Δ 5-desaturase and the deduced amino acid sequence.

Figure 4 shows the deduced amino acid sequence of the PCR fragment (see Example 1)

- 15 Figure 5A and 5B show alignments of the protein sequence of the Δ 5-desaturase with Δ 6-desaturases.

Figure 6A and 6B show the effect of the timing of substrate addition relative to induction on conversion of substrate to product in SC334 containing the Δ 5-desaturase gene.

- 20 Figure 7A and 7B show the effect of inducer concentration on Δ 5-desaturase expression in SC334.

Figure 8A and 8B show the effect of induction temperature on Δ 5-desaturase activity in SC334.

Figure 9A and 9B show the effect of host strain on the conversion of substrate to product in strains expressing the Δ 5-desaturase gene at 15°C.

- 25 Figure 10A and 10B show the effect of host strain on the conversion of substrate to product in strains expressing the Δ 5-desaturase gene at 30°C.

Figure 11 shows the effect of a host strain expressing choline transferase as well as the $\Delta 5$ -desaturase gene on the conversion of substrate to product.

Figure 12A and 12B show the effect of media composition and temperature on the conversion of substrate to product in two host strains
5 expressing the $\Delta 5$ -desaturase gene.

Figure 13 shows alignment of the protein sequence of Ma 29 and contig 253538a.

Figure 14 shows alignment of the protein sequence of Ma 524 and contig 253538a.

10

Brief Description of the Sequence Listings

SEQ ID NO:1 shows a DNA sequence of the *Mortierella alpina* $\Delta 5$ -desaturase.

SEQ ID NO:2 shows an amino acid sequence of *Mortierella alpina* $\Delta 5$ -
15 desaturase.

SEQ ID NO: 3 shows the deduced amino acid sequence of the *M. alpina* PCR fragment (see Example 1).

SEQ ID NO: 4 - SEQ ID NO: 7 show the deduced amino acid sequences of various $\Delta 6$ -desaturases.

20 SEQ ID NO: 8 and SEQ ID NO: 9 show PCR primer sequences for $\Delta 6$ -desaturases

SEQ ID NO: 10 shows a primer for reverse transcription of total RNA.

SEQ ID NO: 11 and SEQ ID NO: 12 show amino acid motifs for desaturase sequences.

25 SEQ ID NO: 13 and SEQ ID NO: 14 show the nucleotide and amino acid sequence of a *Dictyostelium discoideum* desaturase sequence.

SEQ ID NO: 15 and SEQ ID NO: 16 show the nucleotide and amino acid sequence of a *Phaeodactylum tricornutum* desaturase sequence.

SEQ ID NO: 17-20 show the nucleotide and deduced amino acid sequence of a Schizochytrium cDNA clone.

5 SEQ ID NO: 21-27 show nucleotide sequences for human desaturases.

SEQ ID NO: 28 - SEQ ID NO: 34 show peptide sequences for human desaturases.

Detailed Description of the Invention

10 In order to ensure a complete understanding of the invention, the following definitions are provided:

Δ 5-Desaturase: Δ 5 desaturase is an enzyme which introduces a double bond between carbons 5 and 6 from the carboxyl end of a fatty acid molecule.

15 **Δ 6-Desaturase:** Δ 6-desaturase is an enzyme which introduces a double bond between carbons 6 and 7 from the carboxyl end of a fatty acid molecule.

Δ 9-Desaturase: Δ 9-desaturase is an enzyme which introduces a double bond between carbons 9 and 10 from the carboxyl end of a fatty acid molecule.

20 **Δ 12-Desaturase:** Δ 12-desaturase is an enzyme which introduces a double bond between carbons 12 and 13 from the carboxyl end of a fatty acid molecule.

Fatty Acids: Fatty acids are a class of compounds containing a long hydrocarbon chain and a terminal carboxylate group. Fatty acids include the following:

Fatty Acid		
12:0	lauric acid	
16:0	palmitic acid	
16:1	palmitoleic acid	

Fatty Acid		
18:0	stearic acid	
18:1	oleic acid	$\Delta 9$ -18:1
18:2 $\Delta 5,9$	taxoleic acid	$\Delta 5,9$ -18:2
18:2 $\Delta 6,9$	6,9-octadecadienoic acid	$\Delta 6,9$ -18:2
18:2	linoleic acid	$\Delta 9,12$ -18:2 (LA)
18:3 $\Delta 6,9,12$	gamma-linolenic acid	$\Delta 6,9,12$ -18:3 (GLA)
18:3 $\Delta 5,9,12$	pinolenic acid	$\Delta 5,9,12$ -18:3
18:3	alpha-linolenic acid	$\Delta 9,12,15$ -18:3 (ALA)
18:4	stearidonic acid	$\Delta 6,9,12,15$ -18:4 (SDA)
20:0	Arachidic acid	
20:1	Eicosenic Acid	
22:0	behehic acid	
22:1	erucic acid	
22:2	Docasadienoic acid	
20:4 $\omega 6$	arachidonic acid	$\Delta 5,8,11,14$ -20:4 (ARA)
20:3 $\omega 6$	$\omega 6$ -eicosatrienoic dihomo-gamma linolenic	$\Delta 8,11,14$ -20:3 (DGLA)
20:5 $\omega 3$	Eicosapentanoic (Timnodonic acid)	$\Delta 5,8,11,14,17$ -20:5 (EPA)
20:3 $\omega 3$	$\omega 3$ -eicosatrienoic	$\Delta 11,16,17$ -20:3
20:4 $\omega 3$	$\omega 3$ -eicosatetraenoic	$\Delta 8,11,14,17$ -20:4
22:5 $\omega 3$	Docasapentaenoic	$\Delta 7,10,13,16,19$ -22:5 ($\omega 3$ DPA)
22:6 $\omega 3$	Docosaheptaenoic (cervonic acid)	$\Delta 4,7,10,13,16,19$ -22:6 (DHA)
24:0	Lignoceric acid	

Taking into account these definitions, the present invention is directed to novel DNA sequences, DNA constructs, methods and compositions are provided which permit modification of the poly-unsaturated long chain fatty acid content of, for example, microbial cells or animals. Host cells are manipulated to express a sense or antisense transcript of a DNA encoding a polypeptide(s) which catalyzes the conversion of DGLA to ARA. The substrate(s) for the expressed enzyme may be produced by the host cell or may be exogenously supplied. To achieve expression, the transformed DNA is

operably associated with transcriptional and translational initiation and termination regulatory regions that are functional in the host cell. Constructs comprising the gene to be expressed can provide for integration into the genome of the host cell or can autonomously replicate in the host cell. For production of ARA, the expression cassettes generally used include a cassette which provides for $\Delta 5$ -desaturase activity, particularly in a host cell which produces or can take up DGLA. Production of $\omega 6$ -type unsaturated fatty acids, such as ARA, is favored in a host microorganism or animal which is substantially free of ALA. The host is selected or obtained by removing or inhibiting activity of a $\Delta 15$ - or $\omega 3$ - type desaturase (see Figure 2). The endogenous desaturase activity can be affected by providing an expression cassette for an antisense $\Delta 15$ or $\omega 3$ transcript, by disrupting a target $\Delta 15$ - or $\omega 3$ -desaturase gene through insertion, substitution and/or deletion of all or part of the target gene, or by adding a $\Delta 15$ - or $\omega 3$ -desaturase inhibitor. Production of LA also can be increased by providing expression cassettes for $\Delta 9$ and/or $\Delta 12$ -desaturases where their respective enzymatic activities are limiting.

MICROBIAL PRODUCTION OF FATTY ACIDS

Microbial production of fatty acids has several advantages over purification from natural sources such as fish or plants. Many microbes are known with greatly simplified oil compositions compared with those of higher organisms, making purification of desired components easier. Microbial production is not subject to fluctuations caused by external variables such as weather and food supply. Microbially produced oil is substantially free of contamination by environmental pollutants. Additionally, microbes can provide PUFAs in particular forms which may have specific uses. For example, *Spirulina* can provide PUFAs predominantly at the first and third positions of triglycerides; digestion by pancreatic lipases preferentially releases fatty acids from these positions. Following human or animal ingestion of triglycerides derived from *Spirulina*, these PUFAs are released by pancreatic lipases as free

fatty acids and thus are directly available, for example, for infant brain development. Additionally, microbial oil production can be manipulated by controlling culture conditions, notably by providing particular substrates for microbially expressed enzymes, or by addition of compounds which suppress
5 undesired biochemical pathways. In addition to these advantages, production of fatty acids from recombinant microbes provides the ability to alter the naturally occurring microbial fatty acid profile by providing new synthetic pathways in the host or by suppressing undesired pathways, thereby increasing levels of desired PUFAs, or conjugated forms thereof, and decreasing levels of undesired
10 PUFAs.

PRODUCTION OF FATTY ACIDS IN ANIMALS

Production of fatty acids in animals also presents several advantages. Expression of desaturase genes in animals can produce greatly increased levels of desired PUFAs in animal tissues, making recovery from those tissues more
15 economical. For example, where the desired PUFAs are expressed in the breast milk of animals, methods of isolating PUFAs from animal milk are well established. In addition to providing a source for purification of desired PUFAs, animal breast milk can be manipulated through expression of desaturase genes, either alone or in combination with other human genes, to
20 provide animal milks with a PUFA composition substantially similar to human breast milk during the different stages of infant development. Humanized animal milks could serve as infant formulas where human nursing is impossible or undesired, or in cases of malnourishment or disease.

Depending upon the host cell, the availability of substrate, and the
25 desired end product(s), several polypeptides, particularly desaturases, are of interest. By "desaturase" is intended a polypeptide which can desaturate one or more fatty acids to produce a mono- or poly-unsaturated fatty acid or precursor thereof of interest. Of particular interest are polypeptides which can catalyze the conversion of DGLA to produce ARA which includes enzymes which
30 desaturate at the $\Delta 5$ position. By "polypeptide" is meant any chain of amino acids, regardless of length or post-translational modification, for example,

glycosylation or phosphorylation. Considerations for choosing a specific polypeptide having desaturase activity include the pH optimum of the polypeptide, whether the polypeptide is a rate limiting enzyme or a component thereof, whether the desaturase used is essential for synthesis of a desired poly-
5 unsaturated fatty acid, and/or co-factors required by the polypeptide. The expressed polypeptide preferably has parameters compatible with the biochemical environment of its location in the host cell. For example, the polypeptide may have to compete for substrate with other enzymes in the host cell. Analyses of the K_m and specific activity of the polypeptide in question
10 therefore are considered in determining the suitability of a given polypeptide for modifying PUFA production in a given host cell. The polypeptide used in a particular situation is one which can function under the conditions present in the intended host cell but otherwise can be any polypeptide having desaturase activity which has the desired characteristic of being capable of modifying the
15 relative production of a desired PUFA.

For production of ARA, the DNA sequence used encodes a polypeptide having $\Delta 5$ -desaturase activity. In particular instances, this can be coupled with an expression cassette which provides for production of a polypeptide having $\Delta 6$ -desaturase activity and the host cell can optionally be depleted of any $\Delta 15$ -
20 desaturase activity present, for example by providing a transcription cassette for production of antisense sequences to the $\Delta 15$ -desaturase transcription product, by disrupting the $\Delta 15$ -desaturase gene, or by using a host cell which naturally has, or has been mutated to have, low $\Delta 15$ -desaturase activity. Inhibition of undesired desaturase pathways also can be accomplished through the use of
25 specific desaturase inhibitors such as those described in U.S. Patent No. 4,778,630. The choice of combination of cassettes used can depend in part on the PUFA profile of the host cell. Where the host cell $\Delta 5$ -desaturase activity is limiting, overexpression of $\Delta 5$ -desaturase alone generally will be sufficient to provide for enhanced ARA production in the presence of an appropriate
30 substrate such as DGLA. ARA production also can be increased by providing expression cassettes for $\Delta 9$ - or $\Delta 12$ -desaturase genes when the activities of

those desaturases are limiting. A scheme for the synthesis of arachidonic acid (20:4 $\Delta^{5, 8, 11, 14}$) from palmitic acid (C_{16}) is shown in Figure 1. A key enzyme in this pathway is a $\Delta 5$ -desaturase which converts DH- γ -linolenic acid (DGLA, eicosatrienoic acid) to ARA. Conversion of α -linolenic acid (ALA) to stearidonic acid by a $\Delta 6$ -desaturase is also shown. Production of PUFAs in addition to ARA, including EPA and DHA is shown in Figure 2.

SOURCES OF POLYPEPTIDES HAVING DESATURASE ACTIVITY

A source of polypeptides having desaturase activity and oligonucleotides encoding such polypeptides are organisms which produce a desired poly-unsaturated fatty acid. As an example, microorganisms having an ability to produce ARA can be used as a source of $\Delta 5$ -desaturase activity. Such microorganisms include, for example, those belonging to the genera *Mortierella*, *Conidiobolus*, *Pythium*, *Phytophthora*, *Penicillium*, *Porphyridium*, *Coidosporium*, *Mucor*, *Fusarium*, *Aspergillus*, *Rhodotorula*, and *Entomophthora*. Within the genus *Porphyridium*, of particular interest is *Porphyridium cruentum*. Within the genus *Mortierella*, of particular interest are *Mortierella elongata*, *Mortierella exigua*, *Mortierella hygrophila*, *Mortierella ramanniana*, var. *angulispora*, and *Mortierella alpina*. Within the genus *Mucor*, of particular interest are *Mucor circinelloides* and *Mucor javanicus*.

DNAs encoding desired desaturases can be identified in a variety of ways. As an example, a source of the desired desaturase, for example genomic or cDNA libraries from *Mortierella*, is screened with detectable enzymatically- or chemically-synthesized probes, which can be made from DNA, RNA, or non-naturally occurring nucleotides, or mixtures thereof. Probes may be enzymatically synthesized from DNAs of known desaturases for normal or reduced-stringency hybridization methods. Oligonucleotide probes also can be used to screen sources and can be based on sequences of known desaturases, including sequences conserved among known desaturases, or on peptide sequences obtained from the desired purified protein. Oligonucleotide probes based on amino acid sequences can be degenerate to encompass the degeneracy

of the genetic code, or can be biased in favor of the preferred codons of the source organism. Oligonucleotides also can be used as primers for PCR from reverse transcribed mRNA from a known or suspected source; the PCR product can be the full length cDNA or can be used to generate a probe to obtain the
5 desired full length cDNA. Alternatively, a desired protein can be entirely sequenced and total synthesis of a DNA encoding that polypeptide performed.

Once the desired genomic or cDNA has been isolated, it can be sequenced by known methods. It is recognized in the art that such methods are subject to errors, such that multiple sequencing of the same region is routine and
10 is still expected to lead to measurable rates of mistakes in the resulting deduced sequence, particularly in regions having repeated domains, extensive secondary structure, or unusual base compositions, such as regions with high GC base content. When discrepancies arise, resequencing can be done and can employ special methods. Special methods can include altering sequencing conditions
15 by using: different temperatures; different enzymes; proteins which alter the ability of oligonucleotides to form higher order structures; altered nucleotides such as ITP or methylated dGTP; different gel compositions, for example adding formamide; different primers or primers located at different distances from the problem region; or different templates such as single stranded DNAs.
20 Sequencing of mRNA also can be employed.

For the most part, some or all of the coding sequence for the polypeptide having desaturase activity is from a natural source. In some situations, however, it is desirable to modify all or a portion of the codons, for example, to enhance expression, by employing host preferred codons. Host preferred
25 codons can be determined from the codons of highest frequency in the proteins expressed in the largest amount in a particular host species of interest. Thus, the coding sequence for a polypeptide having desaturase activity can be synthesized in whole or in part. All or portions of the DNA also can be synthesized to remove any destabilizing sequences or regions of secondary
30 structure which would be present in the transcribed mRNA. All or portions of the DNA also can be synthesized to alter the base composition to one more

preferable in the desired host cell. Methods for synthesizing sequences and bringing sequences together are well established in the literature. *In vitro* mutagenesis and selection, site-directed mutagenesis, or other means can be employed to obtain mutations of naturally occurring desaturase genes to
5 produce a polypeptide having desaturase activity *in vivo* with more desirable physical and kinetic parameters for function in the host cell, such as a longer half-life or a higher rate of production of a desired polyunsaturated fatty acid.

Mortierella alpina Desaturase

Of particular interest is the *Mortierella alpina* $\Delta 5$ -desaturase which has
10 446 amino acids; the amino acid sequence is shown in Figure 3. The gene encoding the *Mortierella alpina* $\Delta 5$ -desaturase can be expressed in transgenic microorganisms or animals to effect greater synthesis of ARA from DGLA. Other DNAs which are substantially identical to the *Mortierella alpina* $\Delta 5$ -desaturase DNA, or which encode polypeptides which are substantially identical
15 to the *Mortierella alpina* $\Delta 5$ -desaturase polypeptide, also can be used. By substantially identical is intended an amino acid sequence or nucleic acid sequence exhibiting in order of increasing preference at least 60%, 80%, 90% or 95% homology to the *Mortierella alpina* $\Delta 5$ -desaturase amino acid sequence or nucleic acid sequence encoding the amino acid sequence. For polypeptides, the
20 length of comparison sequences generally is at least 16 amino acids, preferably at least 20 amino acids, or most preferably 35 amino acids. For nucleic acids, the length of comparison sequences generally is at least 50 nucleotides, preferably at least 60 nucleotides, and more preferably at least 75 nucleotides, and most preferably, 110 nucleotides. Homology typically is measured using
25 sequence analysis software, for example, the Sequence Analysis software package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wisconsin 53705, MEGAlign (DNASTar, Inc., 1228 S. Park St., Madison, Wisconsin 53715), and MacVector (Oxford Molecular Group, 2105 S. Bascom Avenue, Suite 200,
30 Campbell, California 95008). Such software matches similar sequences by

assigning degrees of homology to various substitutions, deletions, and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine and leucine; aspartic acid, glutamic acid, asparagine, and glutamine; serine and threonine; 5 lysine and arginine; and phenylalanine and tyrosine. Substitutions may also be made on the basis of conserved hydrophobicity or hydrophilicity (Kyte and Doolittle, *J. Mol. Biol.* 157: 105-132, 1982), or on the basis of the ability to assume similar polypeptide secondary structure (Chou and Fasman, *Adv. Enzymol.* 47: 45-148, 1978).

10 Other Desaturases

Encompassed by the present invention are related desaturases from the same or other organisms. Such related desaturases include variants of the disclosed $\Delta 5$ -desaturase naturally occurring within the same or different species of *Mortierella*, as well as homologues of the disclosed $\Delta 5$ -desaturase from other 15 species. Also included are desaturases which, although not substantially identical to the *Mortierella alpina* $\Delta 5$ -desaturase, desaturate a fatty acid molecule at carbon 5 from the carboxyl end of a fatty acid molecule. Related desaturases can be identified by their ability to function substantially the same as the disclosed desaturases; that is, are still able to effectively convert DGLA 20 to ARA. Related desaturases also can be identified by screening sequence databases for sequences homologous to the disclosed desaturase, by hybridization of a probe based on the disclosed desaturase to a library constructed from the source organism, or by RT-PCR using mRNA from the source organism and primers based on the disclosed desaturase. Such 25 desaturases include those from humans, *Dictyostelium discoideum* and *Phaeodactylum tricornum*.

The regions of a desaturase polypeptide important for desaturase activity can be determined through routine mutagenesis, expression of the resulting mutant polypeptides and determination of their activities. Mutants may include 30 deletions, insertions and point mutations, or combinations thereof. A typical

functional analysis begins with deletion mutagenesis to determine the N- and C-terminal limits of the protein necessary for function, and then internal deletions, insertions or point mutants are made to further determine regions necessary for function. Other techniques such as cassette mutagenesis or total synthesis also
5 can be used. Deletion mutagenesis is accomplished, for example, by using exonucleases to sequentially remove the 5' or 3' coding regions. Kits are available for such techniques. After deletion, the coding region is completed by ligating oligonucleotides containing start or stop codons to the deleted coding region after 5' or 3' deletion, respectively. Alternatively, oligonucleotides
10 encoding start or stop codons are inserted into the coding region by a variety of methods including site-directed mutagenesis, mutagenic PCR or by ligation onto DNA digested at existing restriction sites. Internal deletions can similarly be made through a variety of methods including the use of existing restriction sites in the DNA, by use of mutagenic primers via site directed mutagenesis or
15 mutagenic PCR. Insertions are made through methods such as linker-scanning mutagenesis, site-directed mutagenesis or mutagenic PCR. Point mutations are made through techniques such as site-directed mutagenesis or mutagenic PCR.

Chemical mutagenesis also can be used for identifying regions of a desaturase polypeptide important for activity. A mutated construct is expressed,
20 and the ability of the resulting altered protein to function as a desaturase is assayed. Such structure-function analysis can determine which regions may be deleted, which regions tolerate insertions, and which point mutations allow the mutant protein to function in substantially the same way as the native desaturase. All such mutant proteins and nucleotide sequences encoding them
25 are within the scope of the present invention.

EXPRESSION OF DESATURASE GENES

Once the DNA encoding a desaturase polypeptide has been obtained, it is placed in a vector capable of replication in a host cell, or is propagated *in vitro* by means of techniques such as PCR or long PCR. Replicating vectors
30 can include plasmids, phage, viruses, cosmids and the like. Desirable vectors include those useful for mutagenesis of the gene of interest or for expression of

the gene of interest in host cells. The technique of long PCR has made *in vitro* propagation of large constructs possible, so that modifications to the gene of interest, such as mutagenesis or addition of expression signals, and propagation of the resulting constructs can occur entirely *in vitro* without the use of a
5 replicating vector or a host cell.

For expression of a desaturase polypeptide, functional transcriptional and translational initiation and termination regions are operably linked to the DNA encoding the desaturase polypeptide. Expression of the polypeptide coding region can take place *in vitro* or in a host cell. Transcriptional and
10 translational initiation and termination regions are derived from a variety of nonexclusive sources, including the DNA to be expressed, genes known or suspected to be capable of expression in the desired system, expression vectors, chemical synthesis, or from an endogenous locus in a host cell.

Expression In Vitro

15 *In vitro* expression can be accomplished, for example, by placing the coding region for the desaturase polypeptide in an expression vector designed for *in vitro* use and adding rabbit reticulocyte lysate and cofactors; labeled amino acids can be incorporated if desired. Such *in vitro* expression vectors may provide some or all of the expression signals necessary in the system used.
20 These methods are well known in the art and the components of the system are commercially available. The reaction mixture can then be assayed directly for the polypeptide, for example by determining its activity, or the synthesized polypeptide can be purified and then assayed.

Expression In A Host Cell

25 Expression in a host cell can be accomplished in a transient or stable fashion. Transient expression can occur from introduced constructs which contain expression signals functional in the host cell, but which constructs do not replicate and rarely integrate in the host cell, or where the host cell is not proliferating. Transient expression also can be accomplished by inducing the

activity of a regulatable promoter operably linked to the gene of interest, although such inducible systems frequently exhibit a low basal level of expression. Stable expression can be achieved by introduction of a construct that can integrate into the host genome or that autonomously replicates in the host cell. Stable expression of the gene of interest can be selected for through the use of a selectable marker located on or transfected with the expression construct, followed by selection for cells expressing the marker. When stable expression results from integration, integration of constructs can occur randomly within the host genome or can be targeted through the use of constructs containing regions of homology with the host genome sufficient to target recombination with the host locus. Where constructs are targeted to an endogenous locus, all or some of the transcriptional and translational regulatory regions can be provided by the endogenous locus.

When increased expression of the desaturase polypeptide in the source organism is desired, several methods can be employed. Additional genes encoding the desaturase polypeptide can be introduced into the host organism. Expression from the native desaturase locus also can be increased through homologous recombination, for example by inserting a stronger promoter into the host genome to cause increased expression, by removing destabilizing sequences from either the mRNA or the encoded protein by deleting that information from the host genome, or by adding stabilizing sequences to the mRNA (USPN 4,910,141).

When it is desirable to express more than one different gene, appropriate regulatory regions and expression methods, introduced genes can be propagated in the host cell through use of replicating vectors or by integration into the host genome. Where two or more genes are expressed from separate replicating vectors, it is desirable that each vector has a different means of replication. Each introduced construct, whether integrated or not, should have a different means of selection and should lack homology to the other constructs to maintain stable expression and prevent reassortment of elements among constructs. Judicious choices of regulatory regions, selection means and method of

propagation of the introduced construct can be experimentally determined so that all introduced genes are expressed at the necessary levels to provide for synthesis of the desired products.

As an example, where the host cell is a yeast, transcriptional and translational regions functional in yeast cells are provided, particularly from the host species. The transcriptional initiation regulatory regions can be obtained, for example from genes in the glycolytic pathway, such as alcohol dehydrogenase, glyceraldehyde-3-phosphate dehydrogenase (GPD), phosphoglucosomerase, phosphoglycerate kinase, etc. or regulatable genes such as acid phosphatase, lactase, metallothionein, glucoamylase, etc. Any one of a number of regulatory sequences can be used in a particular situation, depending upon whether constitutive or induced transcription is desired, the particular efficiency of the promoter in conjunction with the open-reading frame of interest, the ability to join a strong promoter with a control region from a different promoter which allows for inducible transcription, ease of construction, and the like. Of particular interest are promoters which are activated in the presence of galactose. Galactose-inducible promoters (GAL1, GAL7, and GAL10) have been extensively utilized for high level and regulated expression of protein in yeast (Lue *et al.*, *Mol. Cell. Biol.* Vol. 7, p. 3446, 1987; Johnston, *Microbiol. Rev.* Vol. 51, p. 458, 1987). Transcription from the GAL promoters is activated by the GAL4 protein, which binds to the promoter region and activates transcription when galactose is present. In the absence of galactose, the antagonist GAL80 binds to GAL4 and prevents GAL4 from activating transcription. Addition of galactose prevents GAL80 from inhibiting activation by GAL4.

Nucleotide sequences surrounding the translational initiation codon ATG have been found to affect expression in yeast cells. If the desired polypeptide is poorly expressed in yeast, the nucleotide sequences of exogenous genes can be modified to include an efficient yeast translation initiation sequence to obtain optimal gene expression. For expression in *Saccharomyces*, this can be done by site-directed mutagenesis of an inefficiently expressed gene

by fusing it in-frame to an endogenous *Saccharomyces* gene, preferably a highly expressed gene, such as the lactase gene.

The termination region can be derived from the 3' region of the gene from which the initiation region was obtained or from a different gene. A large number of termination regions are known to and have been found to be satisfactory in a variety of hosts from the same and different genera and species. The termination region usually is selected more as a matter of convenience rather than because of any particular property. Preferably, the termination region is derived from a yeast gene, particularly *Saccharomyces*, *Schizosaccharomyces*, *Candida* or *Kluyveromyces*. The 3' regions of two mammalian genes, γ interferon and $\alpha 2$ interferon, are also known to function in yeast.

INTRODUCTION OF CONSTRUCTS INTO HOST CELLS

Constructs comprising the gene of interest may be introduced into a host cell by standard techniques. These techniques include transformation, protoplast fusion, lipofection, transfection, transduction, conjugation, infection, bolistic impact, electroporation, microinjection, scraping, or any other method which introduces the gene of interest into the host cell. Methods of transformation which are used include lithium acetate transformation (*Methods in Enzymology*, Vol. 194, p. 186-187, 1991). For convenience, a host cell which has been manipulated by any method to take up a DNA sequence or construct will be referred to as "transformed" or "recombinant" herein.

The subject host will have at least have one copy of the expression construct and may have two or more, depending upon whether the gene is integrated into the genome, amplified, or is present on an extrachromosomal element having multiple copy numbers. Where the subject host is a yeast, four principal types of yeast plasmid vectors can be used: Yeast Integrating plasmids (YIps), Yeast Replicating plasmids (YRps), Yeast Centromere plasmids (YCps), and Yeast Episomal plasmids (YEps). YIps lack a yeast replication origin and must be propagated as integrated elements in the yeast genome. YRps have a chromosomally derived autonomously replicating sequence and

are propagated as medium copy number (20 to 40), autonomously replicating, unstably segregating plasmids. YCps have both a replication origin and a centromere sequence and propagate as low copy number (10-20), autonomously replicating, stably segregating plasmids. YEpS have an origin of replication
5 from the yeast 2 μ m plasmid and are propagated as high copy number, autonomously replicating, irregularly segregating plasmids. The presence of the plasmids in yeast can be ensured by maintaining selection for a marker on the plasmid. Of particular interest are the yeast vectors pYES2 (a YEp plasmid available from Invitrogen, confers uracil prototrophy and a GAL1 galactose-
10 inducible promoter for expression), pRS425-pG1 (a YEp plasmid obtained from Dr. T. H. Chang, Ass. Professor of Molecular Genetics, Ohio State University, containing a constitutive GPD promoter and conferring leucine prototrophy), and pYX424 (a YEp plasmid having a constitutive TP1 promoter and conferring leucine prototrophy; Alber, T. and Kawasaki, G. (1982). *J. Mol. & Appl. Genetics* 1: 419).

The transformed host cell can be identified by selection for a marker contained on the introduced construct. Alternatively, a separate marker construct may be introduced with the desired construct, as many transformation techniques introduce many DNA molecules into host cells. Typically,
20 transformed hosts are selected for their ability to grow on selective media. Selective media may incorporate an antibiotic or lack a factor necessary for growth of the untransformed host, such as a nutrient or growth factor. An introduced marker gene therefor may confer antibiotic resistance, or encode an essential growth factor or enzyme, and permit growth on selective media when
25 expressed in the transformed host. Selection of a transformed host also can occur when the expressed marker protein can be detected, either directly or indirectly. The marker protein may be expressed alone or as a fusion to another protein. The marker protein can be detected by its enzymatic activity; for example β galactosidase can convert the substrate X-gal to a colored product,
30 and luciferase can convert luciferin to a light-emitting product. The marker protein can be detected by its light-producing or modifying characteristics; for

example, the green fluorescent protein of *Aequorea victoria* fluoresces when illuminated with blue light. Antibodies can be used to detect the marker protein or a molecular tag on, for example, a protein of interest. Cells expressing the marker protein or tag can be selected, for example, visually, or
5 by techniques such as FACS or panning using antibodies. For selection of yeast transformants, any marker that functions in yeast may be used. Desirably, resistance to kanamycin and the amino glycoside G418 are of interest, as well as ability to grow on media lacking uracil, leucine, lysine or tryptophan.

The $\Delta 5$ -desaturase-mediated production of PUFAs can be performed in
10 either prokaryotic or eukaryotic host cells. Prokaryotic cells of interest include *Eschericia*, *Bacillus*, *Lactobacillus*, *cyanobacteria* and the like. Eukaryotic cells include mammalian cells such as those of lactating animals, avian cells such as of chickens, and other cells amenable to genetic manipulation including insect, fungal, and algae cells. The cells may be cultured or formed as part or
15 all of a host organism including an animal. Viruses and bacteriophage also may be used with the cells in the production of PUFAs, particularly for gene transfer, cellular targeting and selection. In a preferred embodiment, the host is any microorganism or animal which produces DGLA and/or can assimilate exogenously supplied DGLA, and preferably produces large amounts of DGLA.
20 Examples of host animals include mice, rats, rabbits, chickens, quail, turkeys, bovines, sheep, pigs, goats, yaks, etc., which are amenable to genetic manipulation and cloning for rapid expansion of the transgene expressing population. For animals, a $\Delta 5$ -desaturase transgene can be adapted for expression in target organelles, tissues and body fluids through modification of
25 the gene regulatory regions. Of particular interest is the production of PUFAs in the breast milk of the host animal.

Expression In Yeast

Examples of host microorganisms include *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, or other yeast such as *Candida*, *Kluyveromyces*
30 or other fungi, for example, filamentous fungi such as *Aspergillus*, *Neurospora*,

Penicillium, etc. Desirable characteristics of a host microorganism are, for example, that it is genetically well characterized, can be used for high level expression of the product using ultra-high density fermentation, and is on the GRAS (generally recognized as safe) list since the proposed end product is intended for ingestion by humans. Of particular interest is use of a yeast, more particularly baker's yeast (*S. cerevisiae*), as a cell host in the subject invention. Strains of particular interest are SC334 (Mat α pep4-3 prb1-1122 ura3-52 leu2-3, 112 reg1-501 gal1; *Gene* 83:57-64, 1989, Hovland P. *et al.*), YTC34 (α ade2-101 his3 Δ 200 lys2-801 ura3-52; obtained from Dr. T. H. Chang, Ass. Professor of Molecular Genetics, Ohio State University), YTC41 (a/α ura3-52/ura3=52 lys2-801/lys2-801 ade2-101/ade2-101 trp1- Δ 1/trp1- Δ 1 his3 Δ 200/his3 Δ 200 leu2 Δ 1/leu2 Δ 1; obtained from Dr. T. H. Chang, Ass. Professor of Molecular Genetics, Ohio State University), BJ1995 (obtained from the Yeast Genetic Stock Centre, 1021 Donner Laboratory, Berkeley, CA 94720), INVSC1 (Mat α hiw3 Δ 1 leu2 trp1-289 ura3-52; obtained from Invitrogen, 1600 Faraday Ave., Carlsbad, CA 92008) and INVSC2 (Mat α his3 Δ 200 ura3-167; obtained from Invitrogen).

Expression In Avian Species

For producing PUFAs in avian species and cells, such as chickens, turkeys, quail and ducks, gene transfer can be performed by introducing a nucleic acid sequence encoding a Δ 5-desaturase into the cells following procedures known in the art. If a transgenic animal is desired, pluripotent stem cells of embryos can be provided with a vector carrying a Δ 5-desaturase encoding transgene and developed into adult animal (USPN 5,162,215; Ono *et al.* (1996) *Comparative Biochemistry and Physiology A* 113(3):287-292; WO 9612793; WO 9606160). In most cases, the transgene will be modified to express high levels of the desaturase in order to increase production of PUFAs. The transgene can be modified, for example, by providing transcriptional and/or translational regulatory regions that function in avian cells, such as promoters which direct expression in particular tissues and egg parts such as yolk. The

gene regulatory regions can be obtained from a variety of sources, including chicken anemia or avian leukosis viruses or avian genes such as a chicken ovalbumin gene.

Expression In Insect Cells

5 Production of PUFAs in insect cells can be conducted using baculovirus expression vectors harboring a $\Delta 5$ -desaturase transgene. Baculovirus expression vectors are available from several commercial sources such as Clontech. Methods for producing hybrid and transgenic strains of algae, such as marine algae, which contain and express a desaturase transgene also are
10 provided. For example, transgenic marine algae may be prepared as described in USPN 5,426,040. As with the other expression systems described above, the timing, extent of expression and activity of the desaturase transgene can be regulated by fitting the polypeptide coding sequence with the appropriate transcriptional and translational regulatory regions selected for a particular use.
15 Of particular interest are promoter regions which can be induced under preselected growth conditions. For example, introduction of temperature sensitive and/or metabolite responsive mutations into the desaturase transgene coding sequences, its regulatory regions, and/or the genome of cells into which the transgene is introduced can be used for this purpose.

Expression In Plants

20 Production of PUFA's in plants can be conducted using various plant transformation systems such as the use of *Agrobacterium tumefaciens*, plant viruses, particle cell transformation and the like which are disclosed in Applicant's related applications U.S. Application Serial Nos. 08/834,033 and
25 08/956,985 and continuation-in-part applications filed simultaneously with this application all of which are hereby incorporated by reference.

 The transformed host cell is grown under appropriate conditions adapted for a desired end result. For host cells grown in culture, the conditions are

typically optimized to produce the greatest or most economical yield of PUFAs, which relates to the selected desaturase activity. Media conditions which may be optimized include: carbon source, nitrogen source, addition of substrate, final concentration of added substrate, form of substrate added, aerobic or
5 anaerobic growth, growth temperature, inducing agent, induction temperature, growth phase at induction, growth phase at harvest, pH, density, and maintenance of selection. Microorganisms such as yeast, for example, are preferably grown using selected media of interest, which include yeast peptone broth (YPD) and minimal media (contains amino acids, yeast nitrogen base, and
10 ammonium sulfate, and lacks a component for selection, for example uracil). Desirably, substrates to be added are first dissolved in ethanol. Where necessary, expression of the polypeptide of interest may be induced, for example by including or adding galactose to induce expression from a GAL promoter.

15 Expression In An Animal

Expression in cells of a host animal can likewise be accomplished in a transient or stable manner. Transient expression can be accomplished via known methods, for example infection or lipofection, and can be repeated in order to maintain desired expression levels of the introduced construct (*see* Ebert, PCT
20 publication WO 94/05782). Stable expression can be accomplished via integration of a construct into the host genome, resulting in a transgenic animal. The construct can be introduced, for example, by microinjection of the construct into the pronuclei of a fertilized egg, or by transfection, retroviral infection or other techniques whereby the construct is introduced into a cell line which may
25 form or be incorporated into an adult animal (U.S. Patent No. 4,873,191; U.S. Patent No. 5,530,177; U.S. Patent No. 5,565,362; U.S. Patent No. 5,366,894; Wilmut *et al.* (1997) *Nature* 385:810). The recombinant eggs or embryos are transferred to a surrogate mother (U.S. Patent No. 4,873,191; U.S. Patent No. 5,530,177; U.S. Patent No. 5,565,362; U.S. Patent No. 5,366,894; Wilmut *et al.*
30 (*supra*)).

After birth, transgenic animals are identified, for example, by the presence of an introduced marker gene, such as for coat color, or by PCR or Southern blotting from a blood, milk or tissue sample to detect the introduced construct, or by an immunological or enzymological assay to detect the expressed protein or the products produced therefrom (U.S. Patent No. 4,873,191; U.S. Patent No. 5,530,177; U.S. Patent No. 5,565,362; U.S. Patent No. 5,366,894; Wilmut *et al.* (supra)). The resulting transgenic animals may be entirely transgenic or may be mosaics, having the transgenes in only a subset of their cells. The advent of mammalian cloning, accomplished by fusing a nucleated cell with an enucleated egg, followed by transfer into a surrogate mother, presents the possibility of rapid, large-scale production upon obtaining a "founder" animal or cell comprising the introduced construct; prior to this, it was necessary for the transgene to be present in the germ line of the animal for propagation (Wilmut *et al.* (supra)).

Expression in a host animal presents certain efficiencies, particularly where the host is a domesticated animal. For production of PUFAs in a fluid readily obtainable from the host animal, such as milk, the desaturase transgene can be expressed in mammary cells from a female host, and the PUFA content of the host cells altered. The desaturase transgene can be adapted for expression so that it is retained in the mammary cells, or secreted into milk, to form the PUFA reaction products localized to the milk (PCT publication WO 95/24488). Expression can be targeted for expression in mammary tissue using specific regulatory sequences, such as those of bovine α -lactalbumin, α -casein, β -casein, γ -casein, κ -casein, β -lactoglobulin, or whey acidic protein, and may optionally include one or more introns and/or secretory signal sequences (U.S. Patent No. 5,530,177; Rosen, U.S. Patent No. 5,565,362; Clark *et al.*, U.S. Patent No. 5,366,894; Garner *et al.*, PCT publication WO 95/23868). Expression of desaturase transgenes, or antisense desaturase transcripts, adapted in this manner can be used to alter the levels of specific PUFAs, or derivatives thereof, found in the animals milk. Additionally, the $\Delta 5$ -desaturase transgene can be expressed either by itself or with other transgenes, in order to produce

animal milk containing higher proportions of desired PUFAs or PUFA ratios and concentrations that resemble human breast milk (Prieto *et al.*, PCT publication WO 95/24494).

PURIFICATION OF FATTY ACIDS

5 The fatty acids desaturated in the $\Delta 5$ position may be found in the host microorganism or animal as free fatty acids or in conjugated forms such as acylglycerols, phospholipids, sulfolipids or glycolipids, and may be extracted from the host cell through a variety of means well-known in the art. Such means may include extraction with organic solvents, sonication, supercritical
10 fluid extraction using for example carbon dioxide, and physical means such as presses, or combinations thereof. Of particular interest is extraction with methanol and chloroform. Where desirable, the aqueous layer can be acidified to protonate negatively charged moieties and thereby increase partitioning of desired products into the organic layer. After extraction, the organic solvents
15 can be removed by evaporation under a stream of nitrogen. When isolated in conjugated forms, the products may be enzymatically or chemically cleaved to release the free fatty acid or a less complex conjugate of interest, and can then be subject to further manipulations to produce a desired end product. Desirably, conjugated forms of fatty acids are cleaved with potassium hydroxide.

20 If further purification is necessary, standard methods can be employed. Such methods may include extraction, treatment with urea, fractional crystallization, HPLC, fractional distillation, silica gel chromatography, high speed centrifugation or distillation, or combinations of these techniques. Protection of reactive groups, such as the acid or alkenyl groups, may be done at
25 any step through known techniques, for example alkylation or iodination. Methods used include methylation of the fatty acids to produce methyl esters. Similarly, protecting groups may be removed at any step. Desirably, purification of fractions containing ARA, DHA and EPA may be accomplished by treatment with urea and/or fractional distillation.

USES OF FATTY ACIDS

There are several uses for fatty acids of the subject invention. Probes based on the DNAs of the present invention may find use in methods for isolating related molecules or in methods to detect organisms expressing
5 desaturases. When used as probes, the DNAs or oligonucleotides must be detectable. This is usually accomplished by attaching a label either at an internal site, for example via incorporation of a modified residue, or at the 5' or 3' terminus. Such labels can be directly detectable, can bind to a secondary molecule that is detectably labeled, or can bind to an unlabelled secondary
10 molecule and a detectably labeled tertiary molecule; this process can be extended as long as is practical to achieve a satisfactorily detectable signal without unacceptable levels of background signal. Secondary, tertiary, or bridging systems can include use of antibodies directed against any other molecule, including labels or other antibodies, or can involve any molecules
15 which bind to each other, for example a biotin-streptavidin/avidin system. Detectable labels typically include radioactive isotopes, molecules which chemically or enzymatically produce or alter light, enzymes which produce detectable reaction products, magnetic molecules, fluorescent molecules or molecules whose fluorescence or light-emitting characteristics change upon
20 binding. Examples of labelling methods can be found in USPN 5,011,770. Alternatively, the binding of target molecules can be directly detected by measuring the change in heat of solution on binding of probe to target via isothermal titration calorimetry, or by coating the probe or target on a surface and detecting the change in scattering of light from the surface produced by
25 binding of target or probe, respectively, as may be done with the BIAcore system.

PUFAs produced by recombinant means find applications in a wide variety of areas. Supplementation of humans or animals with PUFAs in various forms can result in increased levels not only of the added PUFAs, but of their
30 metabolic progeny as well. For example, where the inherent $\Delta 5$ -desaturase pathway is dysfunctional in an individual, treatment with ARA can result not

only in increased levels of ARA, but also of downstream products of ARA such as prostaglandins (see Figure 1). Complex regulatory mechanisms can make it desirable to combine various PUFAs, or to add different conjugates of PUFAs, in order to prevent, control or overcome such mechanisms to achieve the
5 desired levels of specific PUFAs in an individual.

NUTRITIONAL COMPOSITIONS

The present invention also includes nutritional compositions. Such compositions, for purposes of the present invention, include any food or preparation for human consumption including for enteral or parenteral
10 consumption, which when taken into the body (a) serve to nourish or build up tissues or supply energy and/or (b) maintain, restore or support adequate nutritional status or metabolic function.

The nutritional composition of the present invention comprises at least one oil or acid produced in accordance with the present invention and may
15 either be in a solid or liquid form. Additionally, the composition may include edible macronutrients, vitamins and minerals in amounts desired for a particular use. The amount of such ingredients will vary depending on whether the composition is intended for use with normal, healthy infants, children or adults having specialized needs such as those which accompany certain metabolic
20 conditions (e.g., metabolic disorders).

Examples of macronutrients which may be added to the composition include but are not limited to edible fats, carbohydrates and proteins. Examples of such edible fats include but are not limited to coconut oil, soy oil, and mono- and diglycerides. Examples of such carbohydrates include but are not limited to
25 glucose, edible lactose and hydrolyzed starch. Additionally, examples of proteins which may be utilized in the nutritional composition of the invention include but are not limited to soy proteins, electrodialysed whey , electrodialysed skim milk, milk whey, or the hydrolysates of these proteins.

With respect to vitamins and minerals, the following may be added to
30 the nutritional compositions of the present invention: calcium, phosphorus,

potassium, sodium, chloride, magnesium, manganese, iron, copper, zinc, selenium, iodine, and Vitamins A, E, D, C, and the B complex. Other such vitamins and minerals may also be added.

5 The components utilized in the nutritional compositions of the present invention will of semi-purified or purified origin. By semi-purified or purified is meant a material which has been prepared by purification of a natural material or by synthesis.

Examples of nutritional compositions of the present invention include but are not limited to infant formulas, dietary supplements, and rehydration
10 compositions. Nutritional compositions of particular interest include but are not limited to those utilized for enteral and parenteral supplementation for infants, specialist infant formulae, supplements for the elderly, and supplements for those with gastrointestinal difficulties and/or malabsorption.

Nutritional Compositions

15 A typical nutritional composition of the present invention will contain edible macronutrients, vitamins and minerals in amounts desired for a particular use. The amounts of such ingredients will vary depending on whether the formulation is intended for use with normal, healthy individuals temporarily exposed to stress, or to subjects having specialized needs due to certain chronic
20 or acute disease states (e.g., metabolic disorders). It will be understood by persons skilled in the art that the components utilized in a nutritional formulation of the present invention are of semi-purified or purified origin. By semi-purified or purified is meant a material that has been prepared by purification of a natural material or by synthesis. These techniques are well
25 known in the art (See, e.g., Code of Federal Regulations for Food Ingredients and Food Processing; Recommended Dietary Allowances, 10th Ed., National Academy Press, Washington, D.C., 1989).

In a preferred embodiment, a nutritional formulation of the present invention is an enteral nutritional product, more preferably an adult or child

enteral nutritional product. Accordingly in a further aspect of the invention, a nutritional formulation is provided that is suitable for feeding adults, who are experiencing stress. The formula comprises, in addition to the PUFAs of the invention; macronutrients, vitamins and minerals in amounts designed to
5 provide the daily nutritional requirements of adults.

The macronutritional components include edible fats, carbohydrates and proteins. Exemplary edible fats are coconut oil, soy oil, and mono- and diglycerides and the PUFA oils of this invention. Exemplary carbohydrates are glucose, edible lactose and hydrolyzed cornstarch. A typical protein source
10 would be soy protein, electrodialysed whey or electrodialysed skim milk or milk whey, or the hydrolysates of these proteins, although other protein sources are also available and may be used. These macronutrients would be added in the form of commonly accepted nutritional compounds in amount equivalent to those present in human milk or an energy basis, i.e., on a per calorie basis.

15 Methods for formulating liquid and enteral nutritional formulas are well known in the art and are described in detail in the examples.

The enteral formula can be sterilized and subsequently utilized on a ready-to-feed (RTF) basis or stored in a concentrated liquid or a powder. The powder can be prepared by spray drying the enteral formula prepared as
20 indicated above, and the formula can be reconstituted by rehydrating the concentrate. Adult and infant nutritional formulas are well known in the art and commercially available (e.g., Similac®, Ensure®, Jevity® and Alimentum® from Ross Products Division, Abbott Laboratories). An oil or acid of the present invention can be added to any of these formulas in the amounts
25 described below.

The energy density of the nutritional composition when in liquid form, can typically range from about 0.6 Kcal to 3.0 Kcal per ml. When in solid or powdered form, the nutritional supplement can contain from about 1.2 to more than 9 Kcals per gm, preferably 3 to 7 Kcals per gram. In general, the
30 osmolality of a liquid product should be less than 700 mOsm and more preferably less than 660 mOsm.

The nutritional formula would typically include vitamins and minerals, in addition to the PUFAs of the invention, in order to help the individual ingest the minimum daily requirements for these substances. In addition to the PUFAs listed above, it may also be desirable to supplement the nutritional composition with zinc, copper, and folic acid in addition to antioxidants. It is believed that these substances will also provide a boost to the stressed immune system and thus will provide further benefits to the individual. The presence of zinc, copper or folic acid is optional and is not required in order to gain the beneficial effects on immune suppression. Likewise a pharmaceutical composition can be supplemented with these same substances as well.

In a more preferred embodiment, the nutritional contains, in addition to the antioxidant system and the PUFA component, a source of carbohydrate wherein at least 5 weight % of said carbohydrate is an indigestible oligosaccharide. In yet a more preferred embodiment, the nutritional composition additionally contains protein, taurine and carnitine.

The PUFAs, or derivatives thereof, made by the disclosed method can be used as dietary substitutes, or supplements, particularly infant formulas, for patients undergoing intravenous feeding or for preventing or treating malnutrition. Typically, human breast milk has a fatty acid profile comprising from about 0.15 % to about 0.36 % as DHA, from about 0.03 % to about 0.13 % as EPA, from about 0.30 % to about 0.88 % as ARA, from about 0.22 % to about 0.67 % as DGLA, and from about 0.27 % to about 1.04 % as GLA. Additionally, the predominant triglyceride in human milk has been reported to be 1,3-di-oleoyl-2-palmitoyl, with 2-palmitoyl glycerides reported as better absorbed than 2-oleoyl or 2-lineoyl glycerides (USPN 4,876,107). Thus, fatty acids such as ARA, DGLA, GLA and/or EPA produced by the invention can be used to alter the composition of infant formulas to better replicate the PUFA composition of human breast milk. In particular, an oil composition for use in a pharmacologic or food supplement, particularly a breast milk substitute or supplement, will preferably comprise one or more of ARA, DGLA and GLA.

More preferably the oil will comprise from about 0.3 to 30% ARA, from about 0.2 to 30% DGLA, and from about 0.2 to about 30% GLA.

In addition to the concentration, the ratios of ARA, DGLA and GLA can be adapted for a particular given end use. When formulated as a breast milk supplement, or substitute an oil composition which contains two or more of ARA, DGLA and GLA will be provided in a ratio of about 1:19:30 to about 6:1:0.2, respectively. For example, the breast milk of animals can vary in ratios of ARA:DGLA:DGL ranging from 1:19:30 to 6:1:0.2, which includes intermediate ratios which are preferably about 1:1:1, 1:2:1, 1:1:4. When produced together in a host cell, adjusting the rate and percent of conversion of a precursor substrate such as GLA and DGLA to ARA can be used to precisely control the PUFA ratios. For example, a 5% to 10% conversion rate of DGLA to ARA can be used to produce an ARA to DGLA ratio of about 1:19, whereas a conversion rate of about 75% to 80% can be used to produce an ARA to DGLA ratio of about 6:1. Therefore, whether in a cell culture system or in a host animal, regulating the timing, extent and specificity of desaturase expression as described can be used to modulate the PUFA levels and ratios. Depending on the expression system used, e.g., cell culture or an animal expressing oil(s) in its milk, the oils also can be isolated and recombined in the desired concentrations and ratios. Amounts of oils providing these ratios of PUFA can be determined following standard protocols. PUFAs, or host cells containing them, also can be used as animal food supplements to alter an animal's tissue or milk fatty acid composition to one more desirable for human or animal consumption.

For dietary supplementation, the purified PUFAs, or derivatives thereof, may be incorporated into cooking oils, fats or margarines formulated so that in normal use the recipient would receive the desired amount. The PUFAs may also be incorporated into infant formulas, nutritional supplements or other food products, and may find use as anti-inflammatory or cholesterol lowering agents.

Pharmaceutical Compositions

The present invention also encompasses a pharmaceutical composition comprising one or more of the acids and/or resulting oils produced in accordance with the methods described herein. More specifically, such a
5 pharmaceutical composition may comprise one or more of the acids and/or oils as well as a standard, well-known, non-toxic pharmaceutically acceptable carrier, adjuvant or vehicle such as, for example, phosphate buffered saline, water, ethanol, polyols, vegetable oils, a wetting agent or an emulsion such as a water/oil emulsion. The composition may be in either a liquid or solid form.
10 For example, the composition may be in the form of a tablet, capsule, ingestible liquid or powder, injectible, or topical ointment or cream.

Possible routes of administration include, for example, oral, rectal and parenteral. The route of administration will, of course, depend upon the desired effect. For example, if the composition is being utilized to treat rough, dry, or
15 aging skin, to treat injured or burned skin, or to treat skin or hair affected by a disease or condition, it may perhaps be applied topically.

The dosage of the composition to be administered to the patient may be determined by one of ordinary skill in the art and depends upon various factors such as weight of the patient, age of the patient, immune status of the patient,
20 etc.

With respect to form, the composition may be, for example, a solution, a dispersion, a suspension, an emulsion or a sterile powder which is then reconstituted.

Additionally, the composition of the present invention may be utilized
25 for cosmetic purposes. It may be added to pre-existing cosmetic compositions such that a mixture is formed or may be used as a sole composition.

Pharmaceutical compositions may be utilized to administer the PUFA component to an individual. Suitable pharmaceutical compositions may comprise physiologically acceptable sterile aqueous or non-aqueous solutions,
30 dispersions, suspensions or emulsions and sterile powders for reconstitution into

sterile solutions or dispersions for ingestion. Examples of suitable aqueous and non-aqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglycol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants. It may also be desirable to include isotonic agents, for example sugars, sodium chloride and the like. Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth or mixtures of these substances, and the like.

Solid dosage forms such as tablets and capsules can be prepared using techniques well known in the art. For example, PUFAs of the invention can be tableted with conventional tablet bases such as lactose, sucrose, and cornstarch in combination with binders such as acacia, cornstarch or gelatin, disintegrating agents such as potato starch or alginic acid and a lubricant such as stearic acid or magnesium stearate. Capsules can be prepared by incorporating these excipients into a gelatin capsule along with the antioxidants and the PUFA component. The amount of the antioxidants and PUFA component that should be incorporated into the pharmaceutical formulation should fit within the guidelines discussed above.

As used in this application, the term "treat" refers to either preventing, or reducing the incidence of, the undesired occurrence. For example, to treat immune suppression refers to either preventing the occurrence of this suppression or reducing the amount of such suppression. The terms "patient" and "individual" are being used interchangeably and both refer to an animal. The term "animal" as used in this application refers to any warm-blooded

mammal including, but not limited to, dogs, humans, monkeys, and apes. As used in the application the term "about" refers to an amount varying from the stated range or number by a reasonable amount depending upon the context of use. Any numerical number or range specified in the specification should be
5 considered to be modified by the term about.

"Dose" and "serving" are used interchangeably and refer to the amount of the nutritional or pharmaceutical composition ingested by the patient in a single setting and designed to deliver effective amounts of the antioxidants and the structured triglyceride. As will be readily apparent to those skilled in the
10 art, a single dose or serving of the liquid nutritional powder should supply the amount of antioxidants and PUFAs discussed above. The amount of the dose or serving should be a volume that a typical adult can consume in one sitting. This amount can vary widely depending upon the age, weight, sex or medical condition of the patient. However as a general guideline, a single serving or
15 dose of a liquid nutritional produce should be considered as encompassing a volume from 100 to 600 ml, more preferably from 125 to 500 ml and most preferably from 125 to 300 ml.

The PUFAs of the present invention may also be added to food even when supplementation of the diet is not required. For example, the composition
20 may be added to food of any type including but not limited to margarines, modified butters, cheeses, milk, yogurt, chocolate, candy, snacks, salad oils, cooking oils, cooking fats, meats, fish and beverages.

Pharmaceutical Applications

For pharmaceutical use (human or veterinary), the compositions are
25 generally administered orally but can be administered by any route by which they may be successfully absorbed, e.g., parenterally (i.e. subcutaneously, intramuscularly or intravenously), rectally or vaginally or topically, for example, as a skin ointment or lotion. The PUFAs of the present invention may be administered alone or in combination with a pharmaceutically acceptable
30 carrier or excipient. Where available, gelatin capsules are the preferred form of oral administration. Dietary supplementation as set forth above also can

provide an oral route of administration. The unsaturated acids of the present invention may be administered in conjugated forms, or as salts, esters, amides or prodrugs of the fatty acids. Any pharmaceutically acceptable salt is encompassed by the present invention; especially preferred are the sodium, potassium or lithium salts. Also encompassed are the N-alkylpolyhydroxamine salts, such as N-methyl glucamine, found in PCT publication WO 96/33155. The preferred esters are the ethyl esters. As solid salts, the PUFAs also can be administered in tablet form. For intravenous administration, the PUFAs or derivatives thereof may be incorporated into commercial formulations such as Intralipids. The typical normal adult plasma fatty acid profile comprises 6.64 to 9.46% of ARA, 1.45 to 3.11% of DGLA, and 0.02 to 0.08% of GLA. These PUFAs or their metabolic precursors can be administered, either alone or in mixtures with other PUFAs, to achieve a normal fatty acid profile in a patient. Where desired, the individual components of formulations may be individually provided in kit form, for single or multiple use. A typical dosage of a particular fatty acid is from 0.1 mg to 20 g, or even 100 g daily, and is preferably from 10 mg to 1, 2, 5 or 10 g daily as required, or molar equivalent amounts of derivative forms thereof. Parenteral nutrition compositions comprising from about 2 to about 30 weight percent fatty acids calculated as triglycerides are encompassed by the present invention; preferred is a composition having from about 1 to about 25 weight percent of the total PUFA composition as GLA (USPN 5,196,198). Other vitamins, and particularly fat-soluble vitamins such as vitamin A, D, E and L-carnitine can optionally be included. Where desired, a preservative such as α tocopherol may be added, typically at about 0.1% by weight.

Suitable pharmaceutical compositions may comprise physiologically acceptable sterile aqueous or non-aqueous solutions, dispersions, suspensions or emulsions and sterile powders for reconstitution into sterile injectible solutions or dispersions. Examples of suitable aqueous and non-aqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propyleneglyol, polyethyleneglycol, glycerol, and the like), suitable mixtures thereof, vegetable

oils (such as olive oil) and injectable organic esters such as ethyl oleate. Proper fluidity can be maintained, for example, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants. It may also be desirable to include isotonic agents, for example sugars, sodium chloride and
5 the like. Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents.

Suspensions in addition to the active compounds, may contain suspending agents, as for example, ethoxylated isostearyl alcohols,
10 polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, or mixtures of these substances and the like.

An especially preferred pharmaceutical composition contains diacetyltartaric acid esters of mono- and diglycerides dissolved in an aqueous
15 medium or solvent. Diacetyltartaric acid esters of mono- and diglycerides have an HLB value of about 9-12 and are significantly more hydrophilic than existing antimicrobial lipids that have HLB values of 2-4. Those existing hydrophobic lipids cannot be formulated into aqueous compositions. As disclosed herein, those lipids can now be solubilized into aqueous media in combination with
20 diacetyltartaric acid esters of mono- and diglycerides. In accordance with this embodiment, diacetyltartaric acid esters of mono- and diglycerides (e.g., DATEM-C12:0) is melted with other active antimicrobial lipids (e.g., 18:2 and 12:0 monoglycerides) and mixed to obtain a homogeneous mixture. Homogeneity allows for increased antimicrobial activity. The mixture can be
25 completely dispersed in water. This is not possible without the addition of diacetyltartaric acid esters of mono- and diglycerides and premixing with other monoglycerides prior to introduction into water. The aqueous composition can then be admixed under sterile conditions with physiologically acceptable diluents, preservatives, buffers or propellants as may be required to form a spray
30 or inhalant.

The present invention also encompasses the treatment of numerous disorders with fatty acids. Supplementation with PUFAs of the present invention can be used to treat restenosis after angioplasty. Symptoms of inflammation, rheumatoid arthritis, and asthma and psoriasis can be treated with the PUFAs of the present invention. Evidence indicates that PUFAs may be involved in calcium metabolism, suggesting that PUFAs of the present invention may be used in the treatment or prevention of osteoporosis and of kidney or urinary tract stones.

The PUFAs of the present invention can be used in the treatment of cancer. Malignant cells have been shown to have altered fatty acid compositions; addition of fatty acids has been shown to slow their growth and cause cell death, and to increase their susceptibility to chemotherapeutic agents. GLA has been shown to cause reexpression on cancer cells of the E-cadherin cellular adhesion molecules, loss of which is associated with aggressive metastasis. Clinical testing of intravenous administration of the water soluble lithium salt of GLA to pancreatic cancer patients produced statistically significant increases in their survival. PUFA supplementation may also be useful for treating cachexia associated with cancer.

The PUFAs of the present invention can also be used to treat diabetes (USPN 4,826,877; Horrobin *et al.*, Am. J. Clin. Nutr. Vol. 57 (Suppl.), 732S-737S). Altered fatty acid metabolism and composition has been demonstrated in diabetic animals. These alterations have been suggested to be involved in some of the long-term complications resulting from diabetes, including retinopathy, neuropathy, nephropathy and reproductive system damage. Primrose oil, which contains GLA, has been shown to prevent and reverse diabetic nerve damage.

The PUFAs of the present invention can be used to treat eczema, reduce blood pressure and improve math scores. Essential fatty acid deficiency has been suggested as being involved in eczema, and studies have shown beneficial effects on eczema from treatment with GLA. GLA has also been shown to reduce increases in blood pressure associated with stress, and to improve

performance on arithmetic tests. GLA and DGLA have been shown to inhibit platelet aggregation, cause vasodilation, lower cholesterol levels and inhibit proliferation of vessel wall smooth muscle and fibrous tissue (Brenner *et al.*, Adv. Exp. Med. Biol. Vol. 83, p. 85-101, 1976). Administration of GLA or
5 DGLA, alone or in combination with EPA, has been shown to reduce or prevent gastro-intestinal bleeding and other side effects caused by non-steroidal anti-inflammatory drugs (USPN 4,666,701). GLA and DGLA have also been shown to prevent or treat endometriosis and premenstrual syndrome (USPN 4,758,592) and to treat myalgic encephalomyelitis and chronic fatigue after viral infections
10 (USPN 5,116,871).

Further uses of the PUFAs of this invention include use in treatment of AIDS, multiple sclerosis, acute respiratory syndrome, hypertension and inflammatory skin disorders. The PUFAs of the inventions also can be used for formulas for general health as well as for geriatric treatments.

15

Veterinary Applications

It should be noted that the above-described pharmaceutical and nutritional compositions may be utilized in connection with animals, as well as humans, as animals experience many of the same needs and conditions as human. For example, the oil or acids of the present invention may be utilized in
20 animal feed supplements.

The following examples are presented by way of illustration, not of limitation.

Examples

- | | | |
|----|-----------|------------------------------------------------------------------------------------------|
| 25 | Example 1 | Isolation of a $\Delta 5$ -desaturase Nucleotide Sequence from <i>Mortierella alpina</i> |
| | Example 2 | Expression of <i>M. alpina</i> $\Delta 5$ -desaturase Clones in Baker's Yeast |
| | Example 3 | Initial Optimization of Culture Conditions |

- | | | |
|----|------------|-----------------------------------------------------------------------------------------------------------|
| | Example 4 | Distribution of PUFAs in Yeast Lipid Fractions |
| | Example 5 | Further Culture Optimization |
| | Example 6 | Identification of Homologues to <i>M. alpina</i> $\Delta 5$ and $\Delta 6$ desaturases |
| 5 | Example 7 | Identification of <i>M. alpina</i> $\Delta 5$ and $\Delta 6$ homologues in other PUFA-producing organisms |
| | Example 8 | Identification of <i>M. alpina</i> $\Delta 5$ and $\Delta 6$ homologues in other PUFA-producing organisms |
| | Example 9 | Human Desaturase Sequences |
| 10 | Example 10 | Nutritional Compositions |

Example 1

Isolation of a $\Delta 5$ -desaturase Nucleotide Sequence from *Mortierella alpina*

Mortierella alpina produces arachidonic acid (ARA, 20:4) from the precursor 20:3 by a $\Delta 5$ -desaturase. A nucleotide sequence encoding the $\Delta 5$ -
15 desaturase from *Mortierella alpina* was obtained through PCR amplification using *M. alpina* 1st strand cDNA and degenerate oligonucleotide primers corresponding to amino acid sequences conserved between $\Delta 6$ -desaturases from *Synechocystis* and *Spirulina*. The procedure used was as follows:

Total RNA was isolated from a 3 day old PUFA-producing culture of
20 *Mortierella alpina* using the protocol of Hoge *et al.* (1982) *Experimental Mycology* 6:225-232. The RNA was used to prepare double-stranded cDNA using BRL's lambda-ZipLox system, following the manufacturer's instructions. Several size fractions of the *M. alpina* cDNA were packaged separately to yield libraries with different average-sized inserts. The "full-length" library contains
25 approximately 3×10^6 clones with an average insert size of 1.77 kb. The "sequencing-grade" library contains approximately 6×10^5 clones with an average insert size of 1.1 kb.

5µg of total RNA was reverse transcribed using BRL Superscript RTase and the primer TSyn (5'-CCAAGCTTCTGCAGGAGCTCTTTTTTTT TTTTTTTT-3'), SEQ ID NO:10. Degenerate oligonucleotides were designed to regions conserved between the two cyanobacterial Δ6-desaturase sequences.

5 The specific primers used were D6DESAT-F3 (SEQ ID NO:8) (5'-CUACUACUACUACAYCAYACOTAYACOAAYAT-3') and D6DESAT-R3 (SEQ ID NO:9) (5'-CAUCAUCAUCAUOGGRAAOARRTGRTG-3'), where Y=C+T, R=A+G, and O=I+C. PCR amplification was carried out in a 25µl volume containing: template derived from 40 ng total RNA, 2 pM each primer,

10 200 µM each deoxyribonucleotide triphosphate, 60 mM Tris-Cl, pH 8.5, 15 mM (NH₄)₂SO₄, 2 mM MgCl₂. Samples were subjected to an initial denaturation step of 95 degrees (all temperatures Celsius) for 5 minutes, then held at 72 degrees while 0.2 U of Taq polymerase were added. PCR thermocycling conditions were as follows: 94 degrees for 1 min., 45 degrees for 1.5 min., 72

15 degrees for 2 min. PCR was continued for 35 cycles. PCR using these primers on the *M. alpina* first-strand cDNA produced a 550 bp reaction product. Comparison of the deduced amino acid sequence of the *M. alpina* PCR fragment SEQ ID NO:3 revealed regions of homology with Δ6-desaturases (*see* Figure 5). However, there was only about 28% identity over the region

20 compared.

The PCR product was used as a probe to isolate corresponding cDNA clones from a *M. alpina* library. The longest cDNA clone, Ma29, was designated pCGN5521 and has been completely sequenced on both strands. The cDNA is contained as a 1481 bp insert in the vector pZL1 (Bethesda

25 Research Laboratories) and, beginning with the first ATG, contains an open reading frame encoding 446 amino acids. The reading frame contains the sequence deduced from the PCR fragment. The sequence of the cDNA insert was found to contain regions of homology to Δ6-desaturases (*see* Figure 5). For example, three conserved "histidine boxes" (that have been observed in

30 membrane-bound desaturases (Okuley *et al.*, (1994) *The Plant Cell* 6:147-158)) were found to be present in the *Mortierella* sequence at amino acid positions

171-175, 207-212, and 387-391 (*see* Figure 3). However, the typical "HXXHH" amino acid motif for the third histidine box for the *Mortierella* desaturase was found to be QXXHH, SEQ ID NO:11-12. Surprisingly, the amino-terminus of the encoded protein, showed significant homology to cytochrome b5 proteins. Thus, the *Mortierella* cDNA clone appears to represent a fusion between a cytochrome b5 and a fatty acid desaturase. Since cytochrome b5 is believed to function as the electron donor for membrane-bound desaturase enzymes, it is possible that the N-terminal cytochrome b5 domain of this desaturase protein is involved in its function. This may be advantageous when expressing the desaturase in heterologous systems for PUFA production.

Example 2

Expression of *M. alpina* Desaturase Clones in Baker's Yeast

Yeast Transformation

Lithium acetate transformation of yeast was performed according to standard protocols (*Methods in Enzymology*, Vol. 194, p. 186-187, 1991). Briefly, yeast were grown in YPD at 30°C. Cells were spun down, resuspended in TE, spun down again, resuspended in TE containing 100 mM lithium acetate, spun down again, and resuspended in TE/lithium acetate. The resuspended yeast were incubated at 30°C for 60 minutes with shaking. Carrier DNA was added, and the yeast were aliquoted into tubes. Transforming DNA was added, and the tubes were incubated for 30 min. at 30°C. PEG solution (35% (w/v) PEG 4000, 100 mM lithium acetate, TE pH7.5) was added followed by a 50 min. incubation at 30°C. A 5 min. heat shock at 42°C was performed, the cells were pelleted, washed with TE, pelleted again and resuspended in TE. The resuspended cells were then plated on selective media.

Desaturase Expression in Transformed Yeast

The cDNA clones from *Mortierella alpina* were screened for desaturase activity in baker's yeast. A canola $\Delta 15$ -desaturase (obtained by PCR using 1st

strand cDNA from *Brassica napus* cultivar 212/86 seeds using primers based on the published sequence (Arondel *et al. Science* 258:1353-1355)) was used as a positive control. The $\Delta 15$ -desaturase gene and the gene from cDNA clone Ma29 was inserted into the expression vector pYES2 (Invitrogen), resulting in plasmids pCGR-2 and pCGR-4, respectively. These plasmids were transfected into *S. cerevisiae* yeast strain 334 and expressed after induction with galactose and in the presence of substrates that allowed detection of specific desaturase activity. The control strain was *S. cerevisiae* strain 334 containing the unaltered pYES2 vector. The substrates used, the products produced and the indicated desaturase activity were: DGLA (conversion to ARA would indicate $\Delta 5$ -desaturase activity), linolenic acid (conversion to GLA would indicate $\Delta 6$ -desaturase activity; conversion to ALA would indicate $\Delta 15$ -desaturase activity), oleic acid (an endogenous substrate made by *S. cerevisiae*, conversion to linolenic acid would indicate $\Delta 12$ -desaturase activity, which *S. cerevisiae* lacks), or ARA (conversion to EPA would indicate $\Delta 17$ -desaturase activity). The results are provided in Table 1 below. The lipid fractions were extracted as follows: Cultures were grown for 48-52 hours at 15°C. Cells were pelleted by centrifugation, washed once with sterile ddH₂O, and repelleted. Pellets were vortexed with methanol; chloroform was added along with tritridecanoin (as an internal standard). The mixtures were incubated for at least one hour at room temperature or at 4°C overnight. The chloroform layer was extracted and filtered through a Whatman filter with one gram of anhydrous sodium sulfate to remove particulates and residual water. The organic solvents were evaporated at 40°C under a stream of nitrogen. The extracted lipids were then derivatized to fatty acid methyl esters (FAME) for gas chromatography analysis (GC) by adding 2 ml of 0.5 N potassium hydroxide in methanol to a closed tube. The samples were heated to 95°C to 100°C for 30 minutes and cooled to room temperature. Approximately 2 ml of 14 % boron trifluoride in methanol was added and the heating repeated. After the extracted lipid mixture cooled, 2 ml of water and 1 ml of hexane were added to extract the FAME for analysis by GC. The percent conversion was calculated by dividing the product produced

by the sum of (the product produced and the substrate added) and then multiplying by 100. To calculate the oleic acid percent conversion, as no substrate was added, the total linolenic acid produced was divided by the sum of (oleic acid and linolenic acid produced), then multiplying by 100.

Table 1***M. alpina* Desaturase Expression in Baker's Yeast**

CLONE	TYPE OF ENZYME ACTIVITY	% CONVERSION OF SUBSTRATE
pCGR-2	$\Delta 6$	0 (18:2 to 18:3 $\omega 6$)
(canola $\Delta 15$ desaturase)	$\Delta 15$	16.3 (18:2 to 18:3 $\omega 3$)
	$\Delta 5$	2.0 (20:3 to 20:4 $\omega 6$)
	$\Delta 17$	2.8 (20:4 to 20:5 $\omega 3$)
	$\Delta 12$	1.8 (18:1 to 18:2 $\omega 6$)
pCGR-4	$\Delta 6$	0
(M. alpina Ma29)	$\Delta 15$	0
	$\Delta 5$	15.3
	$\Delta 17$	0.3
	$\Delta 12$	3.3

- 5 The $\Delta 15$ -desaturase control clone exhibited 16.3% conversion of the substrate. The pCGR-4 clone expressing the Ma29 cDNA converted 15.3% of the 20:3 substrate to 20:4 $\omega 6$, indicating that the gene encodes a $\Delta 5$ -desaturase. The background (non-specific conversion of substrate) was between 0-3% in these cases. We also found substrate inhibition of the activity by using
- 10 different concentrations of the substrate. When substrate was added to 100 μM , the percent conversion to product dropped compared to when substrate was added to 25 μM (see below). Additionally, by varying the DGLA substrate concentrations, between about 5 μM to about 200 μM percent conversion of DGLA to ARA ranged from about 5% to 75% with the *M. alpina* $\Delta 5$ -
- 15 desaturase.

These data show that desaturases with different substrate specificities can be expressed in a heterologous system and used to produce poly-unsaturated long chain fatty acids.

Table 2 represents fatty acids of interest as a percent of the total lipid
5 extracted from the yeast host *S. cerevisiae* 334 with the indicated plasmid. No glucose was present in the growth media. Affinity gas chromatography was used to separate the respective lipids. GC/MS was employed to verify the identity product(s). The expected product for the *B. napus* $\Delta 15$ -desaturase, α -
10 linolenic acid, was detected when its substrate, linolenic acid, was added exogenously to the induced yeast culture. This finding demonstrates that yeast expression of a desaturase gene can produce functional enzyme and detectable amounts of product under the current growth conditions. Both exogenously added substrates were taken up by yeast, although slightly less of the longer chain PUFA, dihomo- γ -linolenic acid (20:3), was incorporated into yeast than
15 linolenic acid (18:2) when either was added in free form to the induced yeast cultures. Arachidonic acid was detected as a novel PUFA in yeast when dihomo- γ -linolenic acid was added as the substrate to *S. cerevisiae* 334 (pCGR-4). This identifies pCGR-4 (MA29) as the $\Delta 5$ -desaturase from *M. alpina*. Prior to this, no isolation and expression of a $\Delta 5$ -desaturase from any source has been
20 reported.

Table 2
Fatty Acid as a Percentage of Total Lipid Extracted from Yeast

Plasmid in Yeast (enzyme)	18:2 Incorporated	α -18:3 Produced	γ -18:3 Produced	20:3 Incorporated	20:4 Produced	18:1* Present	18:2 Produced
pYES2 (control)	66.9	0	0	58.4	0	4	0
pCGR-2 (Δ 15)	60.1	5.7	0	50.4	0	0.7	0
pCGR-4 (Δ 5)	67	0	0	32.3	5.8	0.8	0

100 μ M substrate added

* 18:1 is an endogenous fatty acid in yeast

5

- Key To Tables
- 18:1 =oleic acid
 - 18:2 =linolenic acid
 - α -18:3 = α -linolenic acid
 - γ -18:3 = γ -linolenic acid
 - 18:4 =stearidonic acid
 - 20:3 =dihomo- γ -linolenic acid
 - 20:4 =arachidonic acid
- 10

Example 3
Optimization of Culture Conditions

Table 3A shows the effect of exogenous free fatty acid substrate concentration on yeast uptake and conversion to fatty acid product as a percentage of the total yeast lipid extracted. In all instances, low amounts of exogenous substrate (1-10 μM) resulted in low fatty acid substrate uptake and product formation. Between 25 and 50 μM concentration of free fatty acid in the growth and induction media gave the highest percentage of fatty acid product formed, while the 100 μM concentration and subsequent high uptake into yeast appeared to decrease or inhibit the desaturase activity. The feedback inhibition of high fatty acid substrate concentration was well illustrated when the percent conversion rates of the respective fatty acid substrates to their respective products were compared in Table 3B. In all cases, 100 μM substrate concentration in the growth media decreased the percent conversion to product. The effect of media composition was also evident when glucose was present in the growth media for the $\Delta 5$ -desaturase, since the percent of substrate uptake was decreased at 25 μM (Table 3A). However, the percent conversion by $\Delta 5$ -desaturase increased by 18% and the percent product formed remained the same in the presence of glucose in the growth media.

Table 3A
Effect of Added Substrate on the Percentage of Incorporated
Substrate and Product Formed in Yeast Extracts

Plasmid in Yeast	pCGR-2 ($\Delta 15$)	pCGR-4 ($\Delta 5$)
substrate/product	18:2 / α -18:3	20:3/20:4
1 μ M sub.	ND	0.5/1.7
10 μ M sub.	ND	3.3/4
25 μ M sub.	ND	5.1/6.1
25 μ M \diamond sub.	36.6/7.2 \diamond	9.3/5.4 \diamond
50 μ M sub.	53.1/6.5 \diamond	ND
100 μ M sub.	60.1/5.7 \diamond	32.3/5.8 \diamond

5

Table 3B
Effect of Substrate Concentration in Media on the Percent Conversion
of Fatty Acid Substrate to Product in Yeast Extracts

Plasmid in Yeast	pCGR-2 ($\Delta 15$)	pCGR-4 ($\Delta 5$)
substrate/product	18:2 \rightarrow α -18:3	20:3 \rightarrow 20:4
1 μ M sub.	ND	77.3
10 μ M sub.	ND	54.8
25 μ M sub.	ND	54.2
25 μ M \diamond sub.	16.4	36.7
50 μ M sub.	10.9 \diamond	ND
100 μ M sub.	8.7 \diamond	15.2 \diamond

\diamond no glucose in media

* Yeast peptone broth (YPD)

* 18:1 is an endogenous yeast lipid

sub. is substrate concentration

ND (not done)

10

15 Table 4 shows the amount of fatty acid produced by a recombinant desaturase from induced yeast cultures when different amounts of free fatty acid substrate were used. Fatty acid weight was determined since the total amount of lipid varied dramatically when the growth conditions were changed, such as the presence of glucose in the yeast growth and induction media. To better

determine the conditions when the recombinant desaturase would produce the most PUFA product, the quantity of individual fatty acids were examined. The absence of glucose reduced the amount of arachidonic acid produced by $\Delta 5$ -desaturase by half. For $\Delta 5$ -desaturase the amount of total yeast lipid was

5 decreased by almost half in the absence of glucose.

Table 4
Fatty Acid Produced in μg from Yeast Extracts

Plasmid in Yeast (enzyme)	pCGR-4 ($\Delta 5$)	pCGR-7 ($\Delta 12$)
product	20:4	18:2*
1 μM sub.	8.3	ND
10 μM sub.	19.2	ND
25 μM sub.	31.2	115.7
25 μM \diamond sub.	16.8	39 \diamond

\diamond no glucose in media

sub. is substrate concentration

ND (not done)

*18:1, the substrate, is an endogenous yeast lipid

Example 4
Distribution of PUFAs in Yeast Lipid Fractions

15 Table 5 illustrates the uptake of free fatty acids and their new products formed in yeast lipids as distributed in the major lipid fractions. A total lipid extract was prepared as described above. The lipid extract was separated on TLC plates, and the fractions were identified by comparison to standards. The bands were collected by scraping, and internal standards were added. The

20 fractions were then saponified and methylated as above, and subjected to gas chromatography. The gas chromatograph calculated the amount of fatty acid by comparison to a standard. It would appear that the substrates are accessible in the phospholipid form to the desaturases.

Table 5
Fatty Acid Distribution in Various Yeast Lipid Fractions in μg

Fatty acid fraction	Phospholipid	Diglyceride	Free Fatty Acid	Triglyceride	Cholesterol Ester
SC (pCGR-4) substrate 20:3	15.1	1.9	22.9	12.6	3.3
SC (pCGR-4) product 20:4	42.6	0.9	6.8	4.9	0.4

SC = *S. cerevisiae* (plasmid)

5

Example 5

Further Culture Optimization

The growth and induction conditions for optimal activities of desaturases in *Saccharomyces cerevisiae* were evaluated. Various culture conditions that were manipulated for optimal activity were: I) induction temperature, ii) concentration of inducer, iii) timing of substrate addition, iv) concentration of substance, v) sugar source, vi) growth phase at induction. These studies were done using $\Delta 5$ -desaturase gene from *Mortierella alpina* (MA 29). In addition, the effect of changing host strain on expression of the $\Delta 5$ -desaturase gene was also determined.

As described above, the best rate of conversion of substrate to ARA was observed at a substrate concentration of 1 μM , however, the percentage of ARA in the total fatty acids was highest at 25 μM substrate concentration. To determine if the substrate needed to be modified to a readily available form before it could be utilized by the desaturase, the substrate was added either 15 hours before induction or concomitant with inducer addition (indicated as after, in Figure 6A). As it can be seen in Figure 6A, addition of substrate before induction did not have a significant effect on the activity of $\Delta 5$ -desaturase. In fact, addition of substrate along with the inducer was slightly better for expression/activity of $\Delta 5$ -desaturase, as ARA levels in the total fatty acids were

higher. However, the rate of conversion of substrate to product was slightly lower.

The effect of inducer concentration of expression/activity of *Mortierella* $\Delta 5$ -desaturase was examined by inducing SC334/pCGR5 with 0.5 or 2% (w/v) of galactose. As shown in Figures 7A and 7B, expression of $\Delta 5$ -desaturase was higher when induced with 0.5% galactose. Furthermore, rate of conversion of substrate to product was also better when SC334/pCGR5 was induced with 0.5% galactose vs 2% galactose.

To determine the effect of temperature on $\Delta 5$ -desaturase activity, the SC334 host strain, transformed with pCGR5 (SC334/pCGR5) was grown and induced at 15° C, 25° C, 30° C and 37° C. The quantity of ARA (20:4n6) produced in SC334/pCGR5 cultures, supplemental with substrate 20:3n6, was measured by fatty acid analysis. Figure 8A depicts the quantity of 20:3n6 and 20:4n6, expressed as percentage of total fatty acids. Figure 8B depicts the rate of conversion of substrate to product. Growth and induction of SC334/pCGR5 at 25° C, was the best for the expression of $\Delta 5$ -desaturase as evidenced by the highest levels of arachidonic acid in the total fatty acids. Additionally the highest rate of conversion of substrate to product also occurred at 25° C. Growth and induction at 15° C gave the lowest expression of ARA, whereas at 37° C gave the lowest conversion of substrate to product.

The effect of yeast strain on expression of the $\Delta 5$ -desaturase gene was studied in 5 different host strains; INVSC1, INVSC2, YTC34, YTC41, and SC334, at 15° C and 30° C. At 15° C, SC334 has the highest percentage of ARA in total fatty acids, suggesting higher activity of $\Delta 5$ -desaturase in SC334. The rate of conversion of substrate to product, however is lowest in SC334 and highest in INVSC1 (Fig. 9A and B). At 30° C, the highest percentage of product (ARA) in total fatty acids was observed in INVSC2, although the rate of conversion of substrate to product in INVSC2 was slightly lower than INVSC1 (Fig. 10A and B).

ARA, the product of $\Delta 5$ -desaturase, is stored in the phospholipid fraction (Example 4). Therefore the quantity of ARA produced in yeast is limited by the amount that can be stored in the phospholipid fraction. If ARA could also be stored in other fractions such as the triglyceride fraction, the quantity of ARA produced in yeast might be increased. To test this hypothesis, the $\Delta 5$ -desaturase gene was expressed in the yeast host strain DBY746 (obtained from the Yeast Genetic Stock Centre, 1021 Donner Laboratory, Berkeley, CA 94720. The genotype of strain DBY746 is Mat α , his3- $\Delta 1$, leu2-3, leu2-112, ura3-32, trp1-289, gal). The DBY746 yeast strain has an endogenous gene for choline transferase. The presence of this enzyme might enable the DBY746 strain to convert excess phospholipids into triglycerides fraction. Results in Fig. 11 show no increase in the conversion of substrate to product as compared to SC334, which does not have the gene for choline transferase.

To study the effect of media on expression of $\Delta 5$ -desaturase, pCGR4/SC334 was grown in four different media at two different temperatures (15°C and 30°) and in two different host strains (SC334 and INVSC1). The composition of the media was as follows:

Media A: mm-Ura, + 2% galactose + 2% glucose.

Media B: mm-Ura, + 20% galactose + 2% Glucose + 1M sorbitol (pH5.8)

Media C: mm-Ura, + 2% galactose + 2% raffinose

Media D: mm-Ura, + 2% galactose + 2% raffinose + 1M sorbitol (pH5.8)

mm=minimal media

Results show that the highest conversion rate of substrate to product at 15°C in SC334 was observed in media A. The highest conversion rate overall for $\Delta 5$ -desaturase in SC334 was at 30° in media D. The highest conversion rate of $\Delta 5$ -desaturase in INVSC1 was also at 30° in media D (Figures 12A and 12B).

These data show that a DNA encoding a desaturase that can convert DGLA to ARA can be isolated from *Mortierella alpina* and can be expressed in a heterologous system and used to produce poly-unsaturated long chain fatty

acids. Exemplified is the production of ARA from the precursor DGLA by expression of a $\Delta 5$ -desaturase in yeast.

Example 6

Identification of Homologues to *M. alpina* $\Delta 5$ and $\Delta 6$ desaturases

5 A nucleic acid sequence that encodes a putative $\Delta 5$ desaturase was identified through a TBLASTN search of the est databases through NCBI using amino acids 100-446 of Ma29 as a query. The truncated portion of the Ma29 sequence was used to avoid picking up homologies based on the cytochrome b5 portion at the N-terminus of the desaturase. The deduced amino acid sequence
10 of an est from *Dictyostelium discoideum* (accession # C25549) shows very significant homology to Ma29 and lesser, but still significant homology to Ma524. The DNA sequence is presented as SEQ ID NO:13. The amino acid sequence is presented as SEQ ID NO:14.

Example 7

15 **Identification of *M. alpina* $\Delta 5$ and $\Delta 6$ homologues in other**
PUFA-producing organisms

 To look for desaturases involved in PUFA production, a cDNA library was constructed from total RNA isolated from *Phaeodactylum tricornutum*. A plasmid-based cDNA library was constructed in pSPORT1 (GIBCO-BRL)
20 following manufacturer's instructions using a commercially available kit (GIBCO-BRL). Random cDNA clones were sequenced and nucleic acid sequences that encode putative $\Delta 5$ or $\Delta 6$ desaturases were identified through BLAST search of the databases and comparison to Ma29 and Ma524 sequences.

 One clone was identified from the *Phaeodactylum* library with
25 homology to Ma29 and Ma524; it is called 144-011-B12. The DNA sequence is presented as SEQ ID NO:15. The amino acid sequence is presented as SEQ ID NO:16.

Example 8

Identification of *M. alpina* $\Delta 5$ and $\Delta 6$ homologues in other PUFA-producing organisms

To look for desaturases involved in PUFA production, a cDNA library
5 was constructed from total RNA isolated from *Schizochytrium* species. A
plasmid-based cDNA library was constructed in pSPORT1 (GIBCO-BRL)
following manufacturer's instructions using a commercially available kit
(GIBCO-BRL). Random cDNA clones were sequenced and nucleic acid
sequences that encode putative $\Delta 5$ or $\Delta 6$ desaturases were identified through
10 BLAST search of the databases and comparison to Ma29 and Ma524 sequences.

One clone was identified from the *Schizochytrium* library with
homology to Ma29 and Ma524; it is called 81-23-C7. This clone contains a ~1
kb insert. Partial sequence was obtained from each end of the clone using the
universal forward and reverse sequencing primers. The DNA sequence from
15 the forward primer is presented as SEQ ID NO:17. The peptide sequence is
presented as SEQ ID NO:18. The DNA sequence from the reverse primer is
presented as SEQ ID NO:19. The amino acid sequence from the reverse primer
is presented as SEQ ID NO:20.

Example 9

Human Desaturase Gene Sequences

Human desaturase gene sequences potentially involved in long chain
polyunsaturated fatty acid biosynthesis were isolated based on homology
between the human cDNA sequences and *Mortierella alpina* desaturase gene
sequences. The three conserved "histidine boxes" known to be conserved
25 among membrane-bound desaturases were found. As with some other
membrane-bound desaturases the final HXXHH histidine box motif was found
to be QXXHH. The amino acid sequence of the putative human desaturases
exhibited homology to *M. alpina* $\Delta 5$, $\Delta 6$, $\Delta 9$, and $\Delta 12$ desaturases.

The *M. alpina* $\Delta 5$ desaturase and $\Delta 6$ desaturase cDNA sequences were used to search the LifeSeq database of Incyte Pharmaceuticals, Inc., Palo Alto, California 94304. The $\Delta 5$ desaturase sequence was divided into fragments; 1) amino acid no. 1-150, 2) amino acid no. 151-300, and 3) amino acid no. 301-446. The $\Delta 6$ desaturase sequence was divided into three fragments; 1) amino acid no. 1-150, 2) amino acid no. 151-300, and 3) amino acid no. 301-457. These polypeptide fragments were searched against the database using the "tblastn" algorithm. This algorithm compares a protein query sequence against a nucleotide sequence database dynamically translated in all six reading frames (both strands).

The polypeptide fragments 2 and 3 of *M. alpina* $\Delta 5$ and $\Delta 6$ have homologies with the CloneID sequences as outlined in Table 6. The CloneID represents an individual sequence from the Incyte LifeSeq database. After the "tblastn" results have been reviewed, Clone Information was searched with the default settings of Stringency of ≥ 50 , and Productscore ≤ 100 for different CloneID numbers. The Clone Information Results displayed the information including the ClusterID, CloneID, Library, HitID, Hit Description. When selected, the ClusterID number displayed the clone information of all the clones that belong in that ClusterID. The Assemble command assembles all of the CloneID which comprise the ClusterID. The following default settings were used for GCG (Genetics Computer Group, University of Wisconsin Biotechnology Center, Madison, Wisconsin 53705) Assembly:

Word Size:	7
Minimum Overlap:	14
Stringency:	0.8
Minimum Identity:	14
Maximum Gap:	10
Gap Weight:	8
Length Weight:	2

GCG Assembly Results displayed the contigs generated on the basis of sequence information within the CloneID. A contig is an alignment of DNA sequences based on areas of homology among these sequences. A new
5 sequence (consensus sequence) was generated based on the aligned DNA sequences within a contig. The contig containing the CloneID was identified, and the ambiguous sites of the consensus sequence was edited based on the alignment of the CloneIDs (see SEQ ID NO:21 - SEQ ID NO:25) to generate the best possible sequence. The procedure was repeated for all six CloneID
10 listed in Table 6. This produced five unique contigs. The edited consensus sequences of the 5 contigs were imported into the Sequencher software program (Gene Codes Corporation, Ann Arbor, Michigan 48 105). These consensus sequences were assembled. The contig 2511785 overlaps with contig 3506132, and this new contig was called 2535 (SEQ ID NO:27). The contigs from the
15 Sequencher program were copied into the Sequence Analysis software package of GCG.

Each contig was translated in all six reading frames into protein sequences. The *M. alpina* $\Delta 5$ (MA29) and $\Delta 6$ (MA524) sequences were compared with each of the translated contigs using the FastA search (a Pearson
20 and Lipman search for similarity between a query sequence and a group of sequences of the same type (nucleic acid or protein)). Homology among these sequences suggest the open reading frames of each contig. The homology among the *M. alpina* $\Delta 5$ and $\Delta 6$ to contigs 2535 and 3854933 were utilized to create the final contig called 253538a. Figure 13 is the FastA match of the final
25 contig 253538a and MA29, and Figure 14 is the FastA match of the final contig 253538a and MA524. The DNA sequences for the various contigs are presented in SEQ ID NO:21 -SEQ ID NO:27. The various peptide sequences are shown in SEQ ID NO:28 - SEQ ID NO:34.

Although the open reading frame was generated by merging the two
30 contigs, the contig 2535 shows that there is a unique sequence in the beginning of this contig which does not match with the contig 3854933. Therefore, it is

possible that these contigs were generated from independent desaturase like human genes.

The contig 253538a contains an open reading frame encoding 432 amino acids. It starts with Gln (CAG) and ends with the stop codon (TGA).

5 The contig 253538a aligns with both *M. alpina* $\Delta 5$ and $\Delta 6$ sequences, suggesting that it could be either of the desaturases, as well as other known desaturases which share homology with each other. The individual contigs listed in Table 6, as well as the intermediate contig 2535 and the final contig 253538a can be utilized to isolate the complete genes for human desaturases.

10 **Uses of the human desaturases**

These human sequences can be expressed in yeast and plants utilizing the procedures described in the preceding examples. For expression in mammalian cells and transgenic animals, these genes may provide superior codon bias. These human sequences can also be used to identify related

15 desaturase sequences.

Table 6

Sections of the Desaturases	Clone ID from LifeSeq Database	Keyword
151-300 $\Delta 5$	3808675	Fatty acid desaturase
301-446 $\Delta 5$	354535	$\Delta 6$
151-300 $\Delta 6$	3448789	$\Delta 6$
151-300 $\Delta 6$	1362863	$\Delta 6$
151-300 $\Delta 6$	2394760	$\Delta 6$
301-457 $\Delta 6$	3350263	$\Delta 6$

Example 10

Nutritional Compositions

20 The PUFAs of the previous examples can be utilized in various nutritional supplements, infant formulations, nutritional substitutes and other nutrition solutions.

I. INFANT FORMULATIONS

A. Isomil® Soy Formula with Iron.

Usage: As a beverage for infants, children and adults with an allergy or sensitivity to cow's milk. A feeding for patients with disorders for which lactose should be avoided: lactase deficiency, lactose intolerance and galactosemia.

Features:

- Soy protein isolate to avoid symptoms of cow's-milk-protein allergy or sensitivity
- Lactose-free formulation to avoid lactose-associated diarrhea
- Low osmolality (240 mOsm/kg water) to reduce risk of osmotic diarrhea.
- Dual carbohydrates (corn syrup and sucrose) designed to enhance carbohydrate absorption and reduce the risk of exceeding the absorptive capacity of the damaged gut.
- 1.8 mg of Iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.
- Recommended levels of vitamins and minerals.
- Vegetable oils to provide recommended levels of essential fatty acids.
- Milk-white color, milk-like consistency and pleasant aroma.

Ingredients: (Pareve, ©) 85% water, 4.9% corn syrup, 2.6% sugar (sucrose), 2.1% soy oil, 1.9% soy protein isolate, 1.4% coconut oil, 0.15% calcium citrate, 0.11 % calcium phosphate tribasic, potassium citrate, potassium phosphate monobasic, potassium chloride, mono- and diglycerides, soy lecithin, carrageenan, ascorbic acid, L-methionine, magnesium chloride, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic

acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D₃ and cyanocobalamin.

B. Isomil® DF Soy Formula For Diarrhea.

Usage: As a short-term feeding for the dietary management of diarrhea
5 in infants and toddlers.

Features:

- First infant formula to contain added dietary fiber from soy fiber specifically for diarrhea management.
- Clinically shown to reduce the duration of loose, watery stools
10 during mild to severe diarrhea in infants.
- Nutritionally complete to meet the nutritional needs of the infant.
- Soy protein isolate with added L-methionine meets or exceeds an infant's requirement for all essential amino acids.
- Lactose-free formulation to avoid lactose-associated diarrhea.
- 15 • Low osmolality (240 mOsm/kg water) to reduce the risk of osmotic diarrhea.
- Dual carbohydrates (corn syrup and sucrose) designed to enhance carbohydrate absorption and reduce the risk of exceeding the absorptive capacity of the damaged gut.
- 20 • Meets or exceeds the vitamin and mineral levels recommended by the Committee on Nutrition of the American Academy of Pediatrics and required by the Infant Formula Act.
- 1.8 mg of iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.
- 25 • Vegetable oils to provide recommended levels of essential fatty acids.

Ingredients: (Pareve, ☺) 86% water, 4.8% corn syrup, 2.5% sugar (sucrose), 2.1% soy oil, 2.0% soy protein isolate, 1.4% coconut oil, 0.77% soy

fiber, 0.12% calcium citrate, 0.11 % calcium phosphate tribasic, 0.10% potassium citrate, potassium chloride, potassium phosphate monobasic, mono- and diglycerides, soy lecithin, carrageenan, magnesium chloride, ascorbic acid, L-methionine, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D₃ and cyanocobalamin.

10 **C. Isomil® SF Sucrose-Free Soy Formula With Iron.**

Usage: As a beverage for infants, children and adults with an allergy or sensitivity to cow's-milk protein or an intolerance to sucrose. A feeding for patients with disorders for which lactose and sucrose should be avoided.

Features:

- 15 • Soy protein isolate to avoid symptoms of cow's-milk-protein allergy or sensitivity.
- Lactose-free formulation to avoid lactose-associated diarrhea (carbohydrate source is Polycose® Glucose Polymers).
- Sucrose free for the patient who cannot tolerate sucrose.
- 20 • Low osmolality (180 mOsm/kg water) to reduce risk of osmotic diarrhea.
- 1.8 mg of iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.
- Recommended levels of vitamins and minerals.
- 25 • Vegetable oils to provide recommended levels of essential fatty acids.
- Milk-white color, milk-like consistency and pleasant aroma.

Ingredients: (Pareve, ☉) 75% water, 11.8% hydrolyzed cornstarch, 4.1% soy oil, 4.1% soy protein isolate, 2.8% coconut oil, 1.0% modified cornstarch,

0.38% calcium phosphate tribasic, 0.17% potassium citrate, 0.13% potassium chloride, mono- and diglycerides, soy lecithin, magnesium chloride, ascorbic acid, L-methionine, calcium carbonate, sodium chloride, choline chloride, carrageenan, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D₃ and cyanocobalamin.

**D. Isomil® 20 Soy Formula With Iron Ready To Feed,
20 Cal/fl oz.**

Usage: When a soy feeding is desired.

Ingredients: (Pareve, ®) 85% water, 4.9% corn syrup, 2.6% sugar (sucrose), 2.1% soy oil, 1.9% soy protein isolate, 1.4% coconut oil, 0.15% calcium citrate, 0.11% calcium phosphate tribasic, potassium citrate, potassium phosphate monobasic, potassium chloride, mono- and diglycerides, soy lecithin, carrageenan, ascorbic acid, L-methionine, magnesium chloride, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D₃ and cyanocobalamin.

E. Similac® Infant Formula

Usage: When an infant formula is needed: if the decision is made to discontinue breastfeeding before age 1 year, if a supplement to breastfeeding is needed or as a routine feeding if breastfeeding is not adopted.

Features:

- Protein of appropriate quality and quantity for good growth; heat-denatured, which reduces the risk of milk-associated enteric blood loss.

- Fat from a blend of vegetable oils (doubly homogenized), providing essential linoleic acid that is easily absorbed.
- Carbohydrate as lactose in proportion similar to that of human milk.
- Low renal solute load to minimize stress on developing organs.
- Powder, Concentrated Liquid and Ready To Feed forms.

5

Ingredients: (®-D) Water, nonfat milk, lactose, soy oil, coconut oil, mono- and diglycerides, soy lecithin, ascorbic acid, carrageenan, choline chloride, taurine, m-inositol, alpha-tocopheryl acetate, zinc sulfate, niacinamid, ferrous sulfate, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, phyloquinone, biotin, sodium selenite, vitamin D₃ and cyanocobalamin.

10

F. Similac® NeoCare Premature Infant Formula With Iron

Usage: For premature infants' special nutritional needs after hospital discharge. Similac NeoCare is a nutritionally complete formula developed to provide premature infants with extra calories, protein, vitamins and minerals needed to promote catch-up growth and support development.

15

Features:

- Reduces the need for caloric and vitamin supplementation. More calories (22 Cal/fl oz) than standard term formulas (20 Cal/fl oz).
- Highly absorbed fat blend, with medium-chain triglycerides (MCT oil) to help meet the special digestive needs of premature infants.
- Higher levels of protein, vitamins and minerals per 100 Calories to extend the nutritional support initiated in-hospital.
- More calcium and phosphorus for improved bone mineralization.

20

25

Ingredients: ®-D Corn syrup solids, nonfat milk, lactose, whey protein concentrate, soy oil, high-oleic safflower oil, fractionated coconut oil (medium-chain triglycerides), coconut oil, potassium citrate, calcium phosphate tribasic, calcium carbonate, ascorbic acid, magnesium chloride, potassium chloride,

sodium chloride, taurine, ferrous sulfate, m-inositol, choline chloride, ascorbyl palmitate, L-carnitine, alpha-tocopheryl acetate, zinc sulfate, niacinamide, mixed tocopherols, sodium citrate, calcium pantothenate, cupric sulfate, thiamine chloride hydrochloride, vitamin A palmitate, beta carotene, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, phylloquinone, biotin, sodium selenite, vitamin D₃ and cyanocobalamin.

G. Similac Natural Care Low-Iron Human Milk Fortifier Ready To Use, 24 Cal/fl oz.

Usage: Designed to be mixed with human milk or to be fed alternatively with human milk to low-birth-weight infants.

Ingredients: ©-D Water, nonfat milk, hydrolyzed cornstarch, lactose, fractionated coconut oil (medium-chain triglycerides), whey protein concentrate, soil oil, coconut oil, calcium phosphate tribasic, potassium citrate, magnesium chloride, sodium citrate, ascorbic acid, calcium carbonate, mono- and diglycerides, soy lecithin, carrageenan, choline chloride, m-inositol, taurine, niacinamide, L-carnitine, alpha tocopheryl acetate, zinc sulfate, potassium chloride, calcium pantothenate, ferrous sulfate, cupric sulfate, riboflavin, vitamin A palmitate, thiamine chloride hydrochloride, pyridoxine hydrochloride, biotin, folic acid, manganese sulfate, phylloquinone, vitamin D₃, sodium selenite and cyanocobalamin.

Various PUFAs of this invention can be substituted and/or added to the infant formulae described above and to other infant formulae known to those in the art..

II. NUTRITIONAL FORMULATIONS

A. ENSURE®

Usage: ENSURE is a low-residue liquid food designed primarily as an oral nutritional supplement to be used with or between meals or, in appropriate amounts, as a meal replacement. ENSURE is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets. Although it is primarily an oral supplement, it can be fed by tube.

Patient Conditions:

- For patients on modified diets
- For elderly patients at nutrition risk
- For patients with involuntary weight loss
- 5 • For patients recovering from illness or surgery
- For patients who need a low-residue diet

Ingredients:

10 ©-D Water, Sugar (Sucrose), Maltodextrin (Corn), Calcium and Sodium Caseinates, High-Oleic Safflower Oil, Soy Protein Isolate, Soy Oil, Canola Oil, Potassium Citrate, Calcium Phosphate Tribasic, Sodium Citrate, Magnesium Chloride, Magnesium Phosphate Dibasic, Artificial Flavor, Sodium Chloride, Soy Lecithin, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Gellan Gum, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A

15 Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Sodium Molybdate, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate.

B. ENSURE® BARS

20 Usage: ENSURE BARS are complete, balanced nutrition for supplemental use between or with meals. They provide a delicious, nutrient-rich alternative to other snacks. ENSURE BARS contain <1 g lactose/bar, and Chocolate Fudge Brownie flavor is gluten-free. (Honey Graham Crunch flavor contains gluten.)

25 Patient Conditions:

- For patients who need extra calories, protein, vitamins and minerals
- Especially useful for people who do not take in enough calories and nutrients

- For people who have the ability to chew and swallow
- Not to be used by anyone with a peanut allergy or any type of allergy to nuts.

Ingredients:

5 Honey Graham Crunch -- High-Fructose Corn Syrup, Soy Protein Isolate, Brown Sugar, Honey, Maltodextrin (Corn), Crisp Rice (Milled Rice, Sugar [Sucrose], Salt [Sodium Chloride] and Malt), Oat Bran, Partially Hydrogenated Cottonseed and Soy Oils, Soy Polysaccharide, Glycerine, Whey Protein Concentrate, Polydextrose, Fructose, Calcium Caseinate, Cocoa Powder, Artificial Flavors, Canola Oil, High-Oleic Safflower Oil, Nonfat Dry Milk, Whey Powder, Soy Lecithin and Corn Oil. Manufactured in a facility that processes nuts.

Vitamins and Minerals:

15 Calcium Phosphate Tribasic, Potassium Phosphate Dibasic, Magnesium Oxide, Salt (Sodium Chloride), Potassium Chloride, Ascorbic Acid, Ferric Orthophosphate, Alpha-Tocopheryl Acetate, Niacinamide, Zinc Oxide, Calcium Pantothenate, Copper Gluconate, Manganese Sulfate, Riboflavin, Beta-Carotene, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Sodium Molybdate, 20 Phylloquinone, Vitamin D₃ and Cyanocobalamin.

Protein:

Honey Graham Crunch - The protein source is a blend of soy protein isolate and milk proteins.

25	Soy protein isolate	74%
	Milk proteins	26%

Fat:

Honey Graham Crunch - The fat source is a blend of partially hydrogenated cottonseed and soybean, canola, high oleic safflower, and corn oils, and soy lecithin.

	Partially hydrogenated cottonseed and soybean oil	76%
	Canola oil	8%
	High-oleic safflower oil	8%
	Corn oil	4%
5	Soy lecithin	4%

Carbohydrate:

Honey Graham Crunch - The carbohydrate source is a combination of high-fructose corn syrup, brown sugar, maltodextrin, honey, crisp rice, glycerine, soy polysaccharide, and oat bran.

10	High-fructose corn syrup	24%
	Brown sugar	21%
	Maltodextrin	12%
	Honey	11%
	Crisp rice	9%
15	Glycerine	9%
	Soy polysaccharide	7%
	Oat bran	7%\

C. ENSURE® HIGH PROTEIN

20 Usage: ENSURE HIGH PROTEIN is a concentrated, high-protein liquid food designed for people who require additional calories, protein, vitamins, and minerals in their diets. It can be used as an oral nutritional supplement with or between meals or, in appropriate amounts, as a meal replacement. ENSURE HIGH PROTEIN is lactose- and gluten-free, and is
25 suitable for use by people recovering from general surgery or hip fractures and by patients at risk for pressure ulcers.

Patient Conditions

- For patients who require additional calories, protein, vitamins, and minerals, such as patients recovering from general surgery or hip fractures, patients at risk
30 for pressure ulcers, and patients on low-cholesterol diets

Features-

- Low in saturated fat
- Contains 6 g of total fat and < 5 mg of cholesterol per serving
- Rich, creamy taste
- 5 • Excellent source of protein, calcium, and other essential vitamins and minerals
- For low-cholesterol diets
- Lactose-free, easily digested

Ingredients:

10 **Vanilla Supreme:** -D Water, Sugar (Sucrose), Maltodextrin (Corn), Calcium and Sodium Caseinates, High-Oleic Safflower Oil, Soy Protein Isolate, Soy Oil, Canola Oil, Potassium Citrate, Calcium Phosphate Tribasic, Sodium Citrate, Magnesium Chloride, Magnesium Phosphate Dibasic, Artificial Flavor, Sodium Chloride, Soy Lecithin, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc
15 Sulfate, Ferrous Suffate, Alpha-Tocopheryl Acetate, Gellan Gum, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folio Acid, Sodium Motybdate, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D.3 and
20 Cyanocobalarnin.

Protein:

The protein source is a blend of two high-biologic-value proteins: casein and soy.

	Sodium and calcium caseinates	85%
25	Soy protein isolate	15%

Fat:

The fat source is a blend of three oils: high-oleic safflower, canola, and soy.

High-oleic safflower oil	40%
--------------------------	-----

Canola oil	30%
Soy oil	30%

The level of fat in ENSURE HIGH PROTEIN meets American Heart Association (AHA) guidelines. The 6 grams of fat in ENSURE HIGH PROTEIN represent 24% of the total calories, with 2.6% of the fat being from saturated fatty acids and 7.9% from polyunsaturated fatty acids. These values are within the AHA guidelines of $\leq 30\%$ of total calories from fat, $< 10\%$ of the calories from saturated fatty acids, and $\leq 10\%$ of total calories from polyunsaturated fatty acids.

Carbohydrate:

ENSURE HIGH PROTEIN contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla supreme, chocolate royal, wild berry, and banana), plus VARI-FLAVORSO® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla and other nonchocolate flavors

Sucrose	60%
Maltodextrin	40%

Chocolate

Sucrose	70%
Maltodextrin	30%

D. ENSURE® LIGHT

Usage: ENSURE LIGHT is a low-fat liquid food designed for use as an oral nutritional supplement with or between meals. ENSURE LIGHT is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

Patient Conditions:

- For normal-weight or overweight patients who need extra nutrition in a supplement that contains 50% less fat and 20% fewer calories than ENSURE
- For healthy adults who don't eat right and need extra nutrition

Features:

- 5 • Low in fat and saturated fat
- Contains 3 g of total fat per serving and < 5 mg cholesterol
- Rich, creamy taste
- Excellent source of calcium and other essential vitamins and minerals
- For low-cholesterol diets
- 10 • Lactose-free, easily digested

Ingredients:

French Vanilla: ®-D Water, Maltodextrin (Corn), Sugar (Sucrose), Calcium Caseinate, High-Oleic Safflower Oil, Canola Oil, Magnesium Chloride, Sodium Citrate, Potassium Citrate, Potassium Phosphate Dibasic, Magnesium Phosphate Dibasic, Natural and Artificial Flavor, Calcium Phosphate Tribasic, Cellulose Gel, Choline Chloride, Soy Lecithin, Carrageenan, Salt (Sodium Chloride), Ascorbic Acid, Cellulose Gum, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Pyridoxine Hydrochloride, Riboflavin, Chromium Chloride, Folic Acid, Sodium Molybdate, Biotin, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D₃ and Cyanocobalamin.

Protein:

The protein source is calcium caseinate.

25	Calcium caseinate	100%
----	-------------------	------

Fat

The fat source is a blend of two oils: high-oleic safflower and canola.

High-oleic safflower oil	70%
Canola oil	30%

5 The level of fat in ENSURE LIGHT meets American Heart Association (AHA) guidelines. The 3 grams of fat in ENSURE LIGHT represent 13.5% of the total calories, with 1.4% of the fat being from saturated fatty acids and 2.6% from polyunsaturated fatty acids. These values are within the AHA guidelines of $\leq 30\%$ of total calories from fat, $< 10\%$ of the calories from saturated fatty acids, and $\leq 10\%$ of total calories from polyunsaturated fatty acids.

Carbohydrate

10 ENSURE LIGHT contains a combination of maltodextrin and sucrose. The chocolate flavor contains corn syrup as well. The mild sweetness and flavor variety (French vanilla, chocolate supreme, strawberry swirl), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

15 Vanilla and other nonchocolate flavors

Sucrose	51%
Maltodextrin	49%

Chocolate

20	Sucrose	47.0%
	Corn Syrup	26.5%
	Maltodextrin	26.5%

Vitamins and Minerals

An 8-fl-oz serving of ENSURE LIGHT provides at least 25% of the RDIs for 24 key vitamins and minerals.

25 Caffeine

Chocolate flavor contains 2.1 mg caffeine/8 fl oz.

E. ENSURE PLUS®

Usage: ENSURE PLUS is a high-calorie, low-residue liquid food for use when extra calories and nutrients, but a normal concentration of protein, are needed. It is designed primarily as an oral nutritional supplement to be used with or between meals or, in appropriate amounts, as a meal replacement. ENSURE PLUS is lactose- and gluten-free. Although it is primarily an oral nutritional supplement, it can be fed by tube.

Patient Conditions:

- For patients who require extra calories and nutrients, but a normal concentration of protein, in a limited volume
- For patients who need to gain or maintain healthy weight

Features

- Rich, creamy taste
- Good source of essential vitamins and minerals

Ingredients

Vanilla: ®-D Water, Corn Syrup, Maltodextrin (Corn), Corn Oil, Sodium and Calcium Caseinates, Sugar (Sucrose), Soy Protein Isolate, Magnesium Chloride, Potassium Citrate, Calcium Phosphate Tribasic, Soy Lecithin, Natural and Artificial Flavor, Sodium Citrate, Potassium Chloride, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenite, Phylloquinone, Cyanocobalamin and Vitamin D₃.

Protein

The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	84%
-------------------------------	-----

Soy protein isolate	16%
---------------------	-----

Fat

The fat source is corn oil.

Corn oil	100%
----------	------

5 **Carbohydrate**

ENSURE PLUS contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla, chocolate, strawberry, coffee, butter pecan, and eggnog), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

10 **Vanilla, strawberry, butter pecan, and coffee flavors**

Corn Syrup	39%
Maltodextrin	38%
Sucrose	23%

15 **Chocolate and eggnog flavors**

Corn Syrup	36%
Maltodextrin	34%
Sucrose	30%

Vitamins and Minerals

20 An 8-fl-oz serving of ENSURE PLUS provides at least 15% of the RDIs for 25 key Vitamins and minerals.

Caffeine

Chocolate flavor contains 3.1 mg Caffeine/8 fl oz. Coffee flavor contains a trace amount of caffeine.

25

F. ENSURE PLUS® HN

Usage: ENSURE PLUS HN is a nutritionally complete high-calorie, high-nitrogen liquid food designed for people with higher calorie and protein needs or limited volume tolerance. It may be used for oral supplementation or for total nutritional support by tube. ENSURE PLUS HN is lactose- and gluten-free.

Patient Conditions:

- For patients with increased calorie and protein needs, such as following surgery or injury
- For patients with limited volume tolerance and early satiety

Features

- For supplemental or total nutrition
- For oral or tube feeding
- 1.5 CaVmL
- High nitrogen
- Calorically dense

Ingredients

Vanilla: ®-D Water, Maltodextrin (Corn), Sodium and Calcium Caseinates, Corn Oil, Sugar (Sucrose), Soy Protein Isolate, Magnesium Chloride, Potassium Citrate, Calcium Phosphate Tribasic, Soy Lecithin, Natural and Artificial Flavor, Sodium Citrate, Choline Chloride, Ascorbic Acid, Taurine, L-Carnitine, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Carrageenan, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenite, Phylloquinone, Cyanocobalamin and Vitamin D₃.

G. ENSURE® POWDER

Usage: ENSURE POWDER (reconstituted with water) is a low-residue liquid food designed primarily as an oral nutritional supplement to be used with or between meals. ENSURE POWDER is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

5 **Patient Conditions:**

- For patients on modified diets
- For elderly patients at nutrition risk
- For patients recovering from illness/surgery
- For patients who need a low-residue diet

10 **Features**

- Convenient, easy to mix
 - Low in saturated fat
 - Contains 9 g of total fat and < 5 mg of cholesterol per serving
 - High in vitamins and minerals
- 15 • For low-cholesterol diets
- Lactose-free, easily digested

Ingredients: @-D Corn Syrup, Maltodextrin (Corn), Sugar (Sucrose), Corn Oil, Sodium and Calcium Caseinates, Soy Protein Isolate, Artificial Flavor, Potassium Citrate, Magnesium Chloride, Sodium Citrate, Calcium Phosphate

20 Tribasic, Potassium Chloride, Soy Lecithin, Ascorbic Acid, Choline Chloride, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Chloride Hydrochloride, Cupric Sulfate, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Sodium Molybdate, Chromium Chloride, Potassium Iodide,

25 Sodium Selenate, Phylloquinone, Vitamin D₃ and Cyanocobalamin.

Protein

The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	84%
Soy protein isolate	16%

Fat

The fat source is corn oil.

5	Corn oil	100%
---	----------	------

Carbohydrate

ENSURE POWDER contains a combination of corn syrup, maltodextrin, and sucrose. The mild sweetness of ENSURE POWDER, plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, helps to prevent flavor fatigue and aid in patient compliance.

10

Vanilla

Corn Syrup	35%
Maltodextrin	35%
Sucrose	30%

15

H. ENSURE® PUDDING

Usage: ENSURE PUDDING is a nutrient-dense supplement providing balanced nutrition in a nonliquid form to be used with or between meals. It is appropriate for consistency-modified diets (e.g., soft, pureed, or full liquid) or for people with swallowing impairments. ENSURE PUDDING is gluten-free.

20

Patient Conditions:

- For patients on consistency-modified diets (e.g., soft, pureed, or full liquid)
- For patients with swallowing impairments

Features

25

- Rich and creamy, good taste
- Good source of essential vitamins and minerals Convenient-needs no refrigeration

- Gluten-free

Nutrient Profile per 5 oz: Calories 250, Protein 10.9%, Total Fat 34.9%, Carbohydrate 54.2%

Ingredients:

- 5 **Vanilla:** ®-D Nonfat Milk, Water, Sugar (Sucrose), Partially Hydrogenated Soybean Oil, Modified Food Starch, Magnesium Sulfate. Sodium Stearoyl Lactylate, Sodium Phosphate Dibasic, Artificial Flavor, Ascorbic Acid, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Choline Chloride, Niacinamide, Manganese Sulfate, Calcium Pantothenate, FD&C Yellow #5,
- 10 Potassium Citrate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, FD&C Yellow #6, Folic Acid, Biotin, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein

The protein source is nonfat milk.

- | | | |
|----|-------------|------|
| 15 | Nonfat milk | 100% |
|----|-------------|------|

Fat

The fat source is hydrogenated soybean oil.

Hydrogenated soybean oil	100%
--------------------------	------

Carbohydrate

- 20 ENSURE PUDDING contains a combination of sucrose and modified food starch. The mild sweetness and flavor variety (vanilla, chocolate, butterscotch, and tapioca) help prevent flavor fatigue. The product contains 9.2 grams of lactose per serving.

Vanilla and other nonchocolate flavors

- | | | |
|----|----------------------|-----|
| 25 | Sucrose | 56% |
| | Lactose | 27% |
| | Modified food starch | 17% |

Chocolate

Sucrose	58%
Lactose	26%
Modified food starch	16%

5 I. ENSURE® WITH FIBER

Usage: ENSURE WITH FIBER is a fiber-containing, nutritionally complete liquid food designed for people who can benefit from increased dietary fiber and nutrients. ENSURE WITH FIBER is suitable for people who do not require a low-residue diet. It can be fed orally or by tube, and can be used as a nutritional supplement to a regular diet or, in appropriate amounts, as a meal replacement. ENSURE WITH FIBER is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

Patient Conditions

- For patients who can benefit from increased dietary fiber and nutrients

15 Features

- New advanced formula-low in saturated fat, higher in vitamins and minerals
- Contains 6 g of total fat and < 5 mg of cholesterol per serving
- Rich, creamy taste
- Good source of fiber
- Excellent source of essential vitamins and minerals
- For low-cholesterol diets
- Lactose- and gluten-free

Ingredients

Vanilla: ®-D Water, Maltodextrin (Corn), Sugar (Sucrose), Sodium and Calcium Caseinates, Oat Fiber, High-Oleic Safflower Oil, Canola Oil, Soy Protein Isolate, Corn Oil, Soy Fiber, Calcium Phosphate Tribasic, Magnesium Chloride, Potassium Citrate, Cellulose Gel, Soy Lecithin, Potassium Phosphate

5 Dibasic, Sodium Citrate, Natural and Artificial Flavors, Choline Chloride, Magnesium Phosphate, Ascorbic Acid, Cellulose Gum, Potassium Chloride, Carrageenan, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Chromium Chloride, Biotin, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D₃ and Cyanocobalamin.

Protein

10 The protein source is a blend of two high-biologic-value proteins- casein and soy.

Sodium and calcium caseinates	80%
-------------------------------	-----

Soy protein isolate	20%
---------------------	-----

Fat

15 The fat source is a blend of three oils: high-oleic safflower, canola, and corn.

High-oleic safflower oil	40%
--------------------------	-----

Canola oil	40%
------------	-----

Corn oil	20%
----------	-----

20 The level of fat in ENSURE WITH FIBER meets American Heart Association (AHA) guidelines. The 6 grams of fat in ENSURE WITH FIBER represent 22% of the total calories, with 2.01 % of the fat being from saturated fatty acids and 6.7% from polyunsaturated fatty acids. These values are within the AHA guidelines of $\leq 30\%$ of total calories from fat, $< 1\%$ of the calories from saturated fatty acids, and $\leq 1\%$ of total calories from polyunsaturated fatty acids.

25

Carbohydrate

ENSURE WITH FIBER contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla, chocolate, and butter

pecan), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla and other nonchocolate flavors

	Maltodextrin	66%
5	Sucrose	25%
	Oat Fiber	7%
	Soy Fiber	2%

Chocolate

	Maltodextrin	55%
10	Sucrose	36%
	Oat Fiber	7%
	Soy Fiber	2%

Fiber

The fiber blend used in ENSURE WITH FIBER consists of oat fiber and soy polysaccharide. This blend results in approximately 4 grams of total dietary fiber per 8-fl-oz can. The ratio of insoluble to soluble fiber is 95:5.

The various nutritional supplements described above and known to others of skill in the art can be substituted and/or supplemented with the PUFAs of this invention.

20 **J. Oxepa™ Nutritional Product**

Oxepa is low-carbohydrate, calorically dense enteral nutritional product designed for the dietary management of patients with or at risk for ARDS. It has a unique combination of ingredients, including a patented oil blend containing eicosapentaenoic acid (EPA from fish oil), γ -linolenic acid (GLA from borage oil), and elevated antioxidant levels.

Caloric Distribution:

- Caloric density is high at 1.5 Cal/mL (355 Cal/8 fl oz), to minimize the volume required to meet energy needs.
- The distribution of Calories in Oxepa is shown in Table 7.

Table 7. Caloric Distribution of Oxepa			
	per 8 fl oz.	per liter	% of Cal
Calories	355	1,500	---
Fat (g)	22.2	93.7	55.2
Carbohydrate (g)	25	105.5	28.1
Protein (g)	14.8	62.5	16.7
Water (g)	186	785	---

5 **Fat:**

- Oxepa contains 22.2 g of fat per 8-fl oz serving (93.7 g/L).
- The fat source is a oil blend of 31.8% canola oil, 25% medium-chain triglycerides (MCTs), 20% borage oil, 20% fish oil, and 3.2 % soy lecithin. The typical fatty acid profile of Oxepa is shown in Table 8.

- 10
- Oxepa provides a balanced amount of polyunsaturated, monounsaturated, and saturated fatty acids, as shown in Table 10.
 - Medium-chain triglycerides (MCTs) -- 25% of the fat blend -- aid gastric emptying because they are absorbed by the intestinal tract without emulsification by bile acids.

- 15
- The various fatty acid components of Oxepa™ nutritional product can be substituted and/or supplemented with the PUFAs of this invention.

Table 8. Typical Fatty Acid Profile			
	% Total Fatty Acids	g/8 fl oz*	g/L*
Caproic (6:0)	0.2	0.04	0.18
Caprylic (8:0)	14.69	3.1	13.07
Capric (10:0)	11.06	2.33	9.87
Palmitic (16:0)	5.59	1.18	4.98
Palmitoleic (16:1n-7)	1.82	0.38	1.62
Stearic (18:0)	1.84	0.39	1.64

Oleic (18:1n-9)	24.44	5.16	21.75
Linoleic (18:2n-6)	16.28	3.44	14.49
α -Linolenic (18:3n-3)	3.47	0.73	3.09
γ -Linolenic (18:3n-6)	4.82	1.02	4.29
Eicosapentaenoic (20:5n-3)	5.11	1.08	4.55
n-3-Docosapentaenoic (22:5n-3)	0.55	0.12	0.49
Docosahexaenoic (22:6n-3)	2.27	0.48	2.02
Others	7.55	1.52	6.72

* Fatty acids equal approximately 95% of total fat.

Table 9. Fat Profile of Oxepa.	
% of total calories from fat	55.2
Polyunsaturated fatty acids	31.44 g/L
Monounsaturated fatty acids	25.53 g/L
Saturated fatty acids	32.38 g/L
n-6 to n-3 ratio	1.75:1
Cholesterol	9.49 mg/8 fl oz 40.1 mg/L

Carbohydrate:

- The carbohydrate content is 25.0 g per 8-fl-oz serving (105.5 g/L).
- 5 • The carbohydrate sources are 45% maltodextrin (a complex carbohydrate) and 55% sucrose (a simple sugar), both of which are readily digested and absorbed.
- 10 • The high-fat and low-carbohydrate content of Oxepa is designed to minimize carbon dioxide (CO₂) production. High CO₂ levels can complicate weaning in ventilator-dependent patients. The low level of carbohydrate also may be useful for those patients who have developed stress-induced hyperglycemia.
- Oxepa is lactose-free.
- 15 Dietary carbohydrate, the amino acids from protein, and the glycerol moiety of fats can be converted to glucose within the body. Throughout this process, the carbohydrate requirements of glucose-dependent tissues (such as the central nervous system and red blood cells) are met. However, a diet free of

carbohydrates can lead to ketosis, excessive catabolism of tissue protein, and loss of fluid and electrolytes. These effects can be prevented by daily ingestion of 50 to 100 g of digestible carbohydrate, if caloric intake is adequate. The carbohydrate level in Oxepa is also sufficient to minimize gluconeogenesis, if energy needs are being met.

Protein:

- Oxepa contains 14.8 g of protein per 8-fl-oz serving (62.5 g/L).
- The total calorie/nitrogen ratio (150:1) meets the need of stressed patients.
- Oxepa provides enough protein to promote anabolism and the maintenance of lean body mass without precipitating respiratory problems. High protein intakes are a concern in patients with respiratory insufficiency. Although protein has little effect on CO₂ production, a high protein diet will increase ventilatory drive.
- The protein sources of Oxepa are 86.8% sodium caseinate and 13.2% calcium caseinate.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated as incorporated by reference.

The invention now being fully described, it will be apparent to one of ordinary skill in the art that many changes and modifications can be made thereto without departing from the spirit or scope of the appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: KNUTZON, DEBORAH
MURKERJI, PRADIP
HUANG, YUNG-SHENG
THURMOND, JENNIFER
CHAUDHARY, SUNITA
LEONARD, AMANDA

(ii) TITLE OF INVENTION: METHODS AND COMPOSITIONS FOR
SYNTHESIS OF LONG CHAIN POLY-UNSATURATED FATTY ACIDS

(iii) NUMBER OF SEQUENCES: 34

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: LIMBACH & LIMBACH LLP
(B) STREET: 2001 FERRY BUILDING
(C) CITY: SAN FRANCISCO
(D) STATE: CALIFORNIA
(E) COUNTRY: USA
(F) ZIP: 94111

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: MICHAEL R. WARD
(B) REGISTRATION NUMBER: 38,651
(C) REFERENCE/DOCKET NUMBER: CGAB-110

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (415) 433-4150
(B) TELEFAX: (415) 433-8716
(C) TELEX: N/A

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1483 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GCTTCCTCCA GTTCATCCTC CATTTGCGCA CCTGCATTCT TTACGACCGT TAAGCAAGAT 60

5 GGAACGGAC CAAGGAAAAA CCTTCACCTG GGAAGAGCTG GCGGCCCATATA ACACCAAGGA 120
 CGACCTACTC TTGGCCATCC GCGGCAGGGT GTACGATGTC ACAAAGTTCT TGAGCCGCCA 180
 10 TCCTGGTGGA GTGGACACTC TCCTGCTCGG AGCTGGCCGA GATGTTACTC CGGTCTTTGA 240
 GATGTATCAC GCGTTTGGGG CTGCAGATGC CATTATGAAG AAGTACTATG TCGGTACACT 300
 GGTCTCGAAT GAGCTGCCCA TCTTCCCGGA GCCAACGGTG TTCCACAAAA CCATCAAGAC 360
 15 GAGAGTCGAG GGCTACTTTA CGGATCGGAA CATTGATCCC AAGAATAGAC CAGAGATCTG 420
 GGGACGATAC GCTCTTATCT TTGGATCCTT GATCGCTTCC TACTACGCGC AGCTCTTTGT 480
 GCCTTTCGTT GTCGAACGCA CATGGCTTCA GGTGGTGTTC GCAATCATCA TGGGATTTCG 540
 GTGCGCACAA GTCGGACTCA ACCCTCTTCA TGATGCGTCT CACTTTTCAG TGACCCACAA 600
 20 CCCCCTGTC TGGAAGATTC TGGGAGCCAC GCACGACTTT TTCAACGGAG CATCGTACCT 660
 GGTGTGGATG TACCAACATA TGCTCGGCCA TCACCCCTAC ACCAACATTG CTGGAGCAGA 720
 TCCCACGTG TCGACGTCTG AGCCCGATGT TCGTCGTATC AAGCCCAACC AAAAGTGGTT 780
 25 TGTCAACCAC ATCAACCAGC ACATGTTTGT TCCTTTCCTG TACGGACTGC TGGCGTTCAA 840
 GGTGCGCATT CAGGACATCA ACATTTTGTA CTTTGTCAAG ACCAATGACG CTATTCTGTG 900
 CAATCCCATC TCGACATGGC AACTGTGAT GTTCTGGGGC GGCAAGGCTT TCTTTGTCTG 960
 30 GTATCGCCTG ATTGTTCCCC TGCAGTATCT GCCCCTGGGC AAGGTGCTGC TCTTGTTTAC 1020
 GGTGCGGGAC ATGGTGTGCT CTTACTGGCT GCGCTGACC TTCCAGGCGA ACCACGTTGT 1080
 35 TGAGGAAGTT CAGTGGCCGT TGCCTGACGA GAACGGGATC ATCCAAAAGG ACTGGGCAGC 1140
 TATGCAGGTC GAGACTACGC AGGATTACGC ACACGATTCG CACCTCTGGA CCAGCATCAC 1200
 TGGCAGCTTG AACTACCAGG CTGTGCACCA TCTGTTCCCC AACGTGTCGC AGCACCATTA 1260
 40 TCCCAGATATT CTGGCCATCA TCAAGAACAC CTGCAGCGAG TACAAGGTTT CATACCTTGT 1320
 CAAGGATACG TTTTGGCAAG CATTTGCTTC ACATTTGGAG CACTTGCGTG TTCTTGGACT 1380
 45 CCGTCCCAAG GAAGAGTAGA AGAAAAAAG CGCCGAATGA AGTATTGCCC CCTTTTCTC 1440
 CAAGAATGGC AAAAGGAGAT CAAGTGACA TTCTCTATGA AGA 1483

(2) INFORMATION FOR SEQ ID NO:2:

50

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 446 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

55

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

60

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Gly Thr Asp Gln Gly Lys Thr Phe Thr Trp Glu Glu Leu Ala Ala

	1		5		10		15									
	His	Asn	Thr	Lys	Asp	Asp	Leu	Leu	Leu	Ala	Ile	Arg	Gly	Arg	Val	Tyr
				20					25					30		
5	Asp	Val	Thr	Lys	Phe	Leu	Ser	Arg	His	Pro	Gly	Gly	Val	Asp	Thr	Leu
			35					40					45			
	Leu	Leu	Gly	Ala	Gly	Arg	Asp	Val	Thr	Pro	Val	Phe	Glu	Met	Tyr	His
10		50					55					60				
	Ala	Phe	Gly	Ala	Ala	Asp	Ala	Ile	Met	Lys	Lys	Tyr	Tyr	Val	Gly	Thr
	65					70					75				80	
15	Leu	Val	Ser	Asn	Glu	Leu	Pro	Ile	Phe	Pro	Glu	Pro	Thr	Val	Phe	His
				85						90					95	
	Lys	Thr	Ile	Lys	Thr	Arg	Val	Glu	Gly	Tyr	Phe	Thr	Asp	Arg	Asn	Ile
20				100					105					110		
	Asp	Pro	Lys	Asn	Arg	Pro	Glu	Ile	Trp	Gly	Arg	Tyr	Ala	Leu	Ile	Phe
			115					120					125			
25	Gly	Ser	Leu	Ile	Ala	Ser	Tyr	Tyr	Ala	Gln	Leu	Phe	Val	Pro	Phe	Val
		130					135					140				
	Val	Glu	Arg	Thr	Trp	Leu	Gln	Val	Val	Phe	Ala	Ile	Ile	Met	Gly	Phe
	145					150				155					160	
30	Ala	Cys	Ala	Gln	Val	Gly	Leu	Asn	Pro	Leu	His	Asp	Ala	Ser	His	Phe
				165					170						175	
	Ser	Val	Thr	His	Asn	Pro	Thr	Val	Trp	Lys	Ile	Leu	Gly	Ala	Thr	His
35				180					185					190		
	Asp	Phe	Phe	Asn	Gly	Ala	Ser	Tyr	Leu	Val	Trp	Met	Tyr	Gln	His	Met
			195					200					205			
40	Leu	Gly	His	His	Pro	Tyr	Thr	Asn	Ile	Ala	Gly	Ala	Asp	Pro	Asp	Val
		210					215					220				
	Ser	Thr	Ser	Glu	Pro	Asp	Val	Arg	Arg	Ile	Lys	Pro	Asn	Gln	Lys	Trp
	225				230						235				240	
45	Phe	Val	Asn	His	Ile	Asn	Gln	His	Met	Phe	Val	Pro	Phe	Leu	Tyr	Gly
				245					250						255	
	Leu	Leu	Ala	Phe	Lys	Val	Arg	Ile	Gln	Asp	Ile	Asn	Ile	Leu	Tyr	Phe
50				260					265					270		
	Val	Lys	Thr	Asn	Asp	Ala	Ile	Arg	Val	Asn	Pro	Ile	Ser	Thr	Trp	His
			275					280					285			
55	Thr	Val	Met	Phe	Trp	Gly	Gly	Lys	Ala	Phe	Phe	Val	Trp	Tyr	Arg	Leu
		290				295						300				
	Ile	Val	Pro	Leu	Gln	Tyr	Leu	Pro	Leu	Gly	Lys	Val	Leu	Leu	Leu	Phe
	305				310						315				320	
60	Thr	Val	Ala	Asp	Met	Val	Ser	Ser	Tyr	Trp	Leu	Ala	Leu	Thr	Phe	Gln
				325						330					335	

	Ala	Asn	His	Val	Val	Glu	Glu	Val	Gln	Trp	Pro	Leu	Pro	Asp	Glu	Asn
				340					345					350		
5	Gly	Ile	Ile	Gln	Lys	Asp	Trp	Ala	Ala	Met	Gln	Val	Glu	Thr	Thr	Gln
			355					360					365			
	Asp	Tyr	Ala	His	Asp	Ser	His	Leu	Trp	Thr	Ser	Ile	Thr	Gly	Ser	Leu
		370					375					380				
10	Asn	Tyr	Gln	Ala	Val	His	His	Leu	Phe	Pro	Asn	Val	Ser	Gln	His	His
	385					390					395					400
	Tyr	Pro	Asp	Ile	Leu	Ala	Ile	Ile	Lys	Asn	Thr	Cys	Ser	Glu	Tyr	Lys
15				405					410						415	
	Val	Pro	Tyr	Leu	Val	Lys	Asp	Thr	Phe	Trp	Gln	Ala	Phe	Ala	Ser	His
				420					425				430			
	Leu	Glu	His	Leu	Arg	Val	Leu	Gly	Leu	Arg	Pro	Lys	Glu	Glu		
20			435					440					445			

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 186 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: not relevant
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

35	Leu	His	His	Thr	Tyr	Asn	Ile	Ala	Gly	Ala	Asp	Pro	Asp	Val	Ser	
	1				5				10					15		
	Thr	Ser	Glu	Pro	Asp	Val	Arg	Arg	Ile	Lys	Pro	Asn	Gln	Lys	Trp	Phe
40				20					25					30		
	Val	Asn	His	Ile	Asn	Gln	His	Met	Phe	Val	Pro	Phe	Leu	Tyr	Gly	Leu
			35					40					45			
45	Leu	Ala	Phe	Lys	Val	Arg	Ile	Gln	Asp	Ile	Asn	Ile	Leu	Tyr	Phe	Val
	50						55					60				
	Lys	Thr	Asn	Asp	Ala	Ile	Arg	Val	Asn	Pro	Ile	Ser	Thr	Trp	His	Thr
	65					70					75					80
50	Val	Met	Phe	Trp	Gly	Gly	Lys	Ala	Phe	Phe	Val	Trp	Tyr	Arg	Leu	Ile
					85					90					95	
	Val	Pro	Leu	Gln	Tyr	Leu	Pro	Leu	Gly	Lys	Val	Leu	Leu	Leu	Phe	Thr
55				100					105					110		
	Val	Ala	Asp	Met	Val	Ser	Ser	Tyr	Trp	Leu	Ala	Leu	Thr	Phe	Gln	Ala
			115					120					125			
60	Asn	Tyr	Val	Val	Glu	Glu	Val	Gln	Trp	Pro	Leu	Pro	Asp	Glu	Asn	Gly
	130						135					140				
	Ile	Ile	Gln	Lys	Asp	Trp	Ala	Ala	Met	Gln	Val	Glu	Thr	Thr	Gln	Asp

```

145                               150                               155                               160
Tyr Ala His Asp Ser His Leu Trp Thr Ser Ile Thr Gly Ser Leu Asn
                    165                               170                               175

5
Tyr Gln Xaa Val His His Leu Phe Pro His
                    180                               185

10
(2) INFORMATION FOR SEQ ID NO:4:

    (i) SEQUENCE CHARACTERISTICS:
        (A) LENGTH: 457 amino acids
        (B) TYPE: amino acid
        (C) STRANDEDNESS: not relevant
        (D) TOPOLOGY: linear

    (ii) MOLECULE TYPE: peptide

20
    (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Ala Ala Ala Pro Ser Val Arg Thr Phe Thr Arg Ala Glu Val Leu
1          5          10          15

25
Asn Ala Glu Ala Leu Asn Glu Gly Lys Lys Asp Ala Glu Ala Pro Phe
          20          25          30

Leu Met Ile Ile Asp Asn Lys Val Tyr Asp Val Arg Glu Phe Val Pro
          35          40          45

30
Asp His Pro Gly Gly Ser Val Ile Leu Thr His Val Gly Lys Asp Gly
          50          55          60

Thr Asp Val Phe Asp Thr Phe His Pro Glu Ala Ala Trp Glu Thr Leu
35          65          70          75          80

Ala Asn Phe Tyr Val Gly Asp Ile Asp Glu Ser Asp Arg Asp Ile Lys
          85          90          95

40
Asn Asp Asp Phe Ala Ala Glu Val Arg Lys Leu Arg Thr Leu Phe Gln
          100          105          110

Ser Leu Gly Tyr Tyr Asp Ser Ser Lys Ala Tyr Tyr Ala Phe Lys Val
          115          120          125

45
Ser Phe Asn Leu Cys Ile Trp Gly Leu Ser Thr Val Ile Val Ala Lys
          130          135          140

Trp Gly Gln Thr Ser Thr Leu Ala Asn Val Leu Ser Ala Ala Leu Leu
50          145          150          155          160

Gly Leu Phe Trp Gln Gln Cys Gly Trp Leu Ala His Asp Phe Leu His
          165          170          175

55
His Gln Val Phe Gln Asp Arg Phe Trp Gly Asp Leu Phe Gly Ala Phe
          180          185          190

Leu Gly Gly Val Cys Gln Gly Phe Ser Ser Ser Trp Trp Lys Asp Lys
          195          200          205

60
His Asn Thr His His Ala Ala Pro Asn Val His Val Glu Asp Pro Asp
          210          215          220

```

Ile Asp Thr His Pro Leu Leu Thr Trp Ser Glu His Ala Leu Glu Met
 225 230 235 240
 5 Phe Ser Asp Val Pro Asp Glu Glu Leu Thr Arg Met Trp Ser Arg Phe
 245 250 255
 Met Val Leu Asn Gln Thr Trp Phe Tyr Phe Pro Ile Leu Ser Phe Ala
 260 265 270
 10 Arg Leu Ser Trp Cys Leu Gln Ser Ile Leu Phe Val Leu Pro Asn Gly
 275 280 285
 Gln Ala His Lys Pro Ser Gly Ala Arg Val Pro Ile Ser Leu Val Glu
 290 295 300
 15 Gln Leu Ser Leu Ala Met His Trp Thr Trp Tyr Leu Ala Thr Met Phe
 305 310 315 320
 20 Leu Phe Ile Lys Asp Pro Val Asn Met Leu Val Tyr Phe Leu Val Ser
 325 330 335
 Gln Ala Val Cys Gly Asn Leu Leu Ala Ile Val Phe Ser Leu Asn His
 340 345 350
 25 Asn Gly Met Pro Val Ile Ser Lys Glu Glu Ala Val Asp Met Asp Phe
 355 360 365
 Phe Thr Lys Gln Ile Ile Thr Gly Arg Asp Val His Pro Gly Leu Phe
 370 375 380
 30 Ala Asn Trp Phe Thr Gly Gly Leu Asn Tyr Gln Ile Glu His His Leu
 385 390 395 400
 35 Phe Pro Ser Met Pro Arg His Asn Phe Ser Lys Ile Gln Pro Ala Val
 405 410 415
 Glu Thr Leu Cys Lys Lys Tyr Asn Val Arg Tyr His Thr Thr Gly Met
 420 425 430
 40 Ile Glu Gly Thr Ala Glu Val Phe Ser Arg Leu Asn Glu Val Ser Lys
 435 440 445
 45 Ala Ala Ser Lys Met Gly Lys Ala Gln
 450 455

(2) INFORMATION FOR SEQ ID NO:5:

- 50 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 446 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met Ala Ala Gln Ile Lys Lys Tyr Ile Thr Ser Asp Glu Leu Lys Asn
 1 5 10 15

His Asp Lys Pro Gly Asp Leu Trp Ile Ser Ile Gln Gly Lys Ala Tyr
 20 25 30
 5 Asp Val Ser Asp Trp Val Lys Asp His Pro Gly Gly Ser Phe Pro Leu
 35 40 45
 Lys Ser Leu Ala Gly Gln Glu Val Thr Asp Ala Phe Val Ala Phe His
 50 55 60
 10 Pro Ala Ser Thr Trp Lys Asn Leu Asp Lys Phe Phe Thr Gly Tyr Tyr
 65 70 75 80
 Leu Lys Asp Tyr Ser Val Ser Glu Val Ser Lys Val Tyr Arg Lys Leu
 85 90 95
 Val Phe Glu Phe Ser Lys Met Gly Leu Tyr Asp Lys Lys Gly His Ile
 100 105 110
 20 Met Phe Ala Thr Leu Cys Phe Ile Ala Met Leu Phe Ala Met Ser Val
 115 120 125
 Tyr Gly Val Leu Phe Cys Glu Gly Val Leu Val His Leu Phe Ser Gly
 130 135 140
 25 Cys Leu Met Gly Phe Leu Trp Ile Gln Ser Gly Trp Ile Gly His Asp
 145 150 155 160
 Ala Gly His Tyr Met Val Val Ser Asp Ser Arg Leu Asn Lys Phe Met
 165 170 175
 30 Gly Ile Phe Ala Ala Asn Cys Leu Ser Gly Ile Ser Ile Gly Trp Trp
 180 185 190
 35 Lys Trp Asn His Asn Ala His His Ile Ala Cys Asn Ser Leu Glu Tyr
 195 200 205
 Asp Pro Asp Leu Gln Tyr Ile Pro Phe Leu Val Val Ser Ser Lys Phe
 210 215 220
 40 Phe Gly Ser Leu Thr Ser His Phe Tyr Glu Lys Arg Leu Thr Phe Asp
 225 230 235 240
 Ser Leu Ser Arg Phe Phe Val Ser Tyr Gln His Trp Thr Phe Tyr Pro
 245 250 255
 45 Ile Met Cys Ala Ala Arg Leu Asn Met Tyr Val Gln Ser Leu Ile Met
 260 265 270
 50 Leu Leu Thr Lys Arg Asn Val Ser Tyr Arg Ala Gln Glu Leu Leu Gly
 275 280 285
 Cys Leu Val Phe Ser Ile Trp Tyr Pro Leu Leu Val Ser Cys Leu Pro
 290 295 300
 55 Asn Trp Gly Glu Arg Ile Met Phe Val Ile Ala Ser Leu Ser Val Thr
 305 310 315 320
 60 Gly Met Gln Gln Val Gln Phe Ser Leu Asn His Phe Ser Ser Ser Val
 325 330 335
 Tyr Val Gly Lys Pro Lys Gly Asn Asn Trp Phe Glu Lys Gln Thr Asp

340 345 350
 Gly Thr Leu Asp Ile Ser Cys Pro Pro Trp Met Asp Trp Phe His Gly
 355 360 365
 5
 Gly Leu Gln Phe Gln Ile Glu His His Leu Phe Pro Lys Met Pro Arg
 370 375 380
 10
 Cys Asn Leu Arg Lys Ile Ser Pro Tyr Val Ile Glu Leu Cys Lys Lys
 385 390 395 400
 His Asn Leu Pro Tyr Asn Tyr Ala Ser Phe Ser Lys Ala Asn Glu Met
 405 410 415
 15
 Thr Leu Arg Thr Leu Arg Asn Thr Ala Leu Gln Ala Arg Asp Ile Thr
 420 425 430
 Lys Pro Leu Pro Lys Asn Leu Val Trp Glu Ala Leu His Thr
 435 440 445
 20

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 359 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear
 25

(ii) MOLECULE TYPE: peptide
30

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

35 Met Leu Thr Ala Glu Arg Ile Lys Phe Thr Gln Lys Arg Gly Phe Arg
 1 5 10 15
 Arg Val Leu Asn Gln Arg Val Asp Ala Tyr Phe Ala Glu His Gly Leu
 20 25 30
 40 Thr Gln Arg Asp Asn Pro Ser Met Tyr Leu Lys Thr Leu Ile Ile Val
 35 40 45
 Leu Trp Leu Phe Ser Ala Trp Ala Phe Val Leu Phe Ala Pro Val Ile
 50 55 60
 45 Phe Pro Val Arg Leu Leu Gly Cys Met Val Leu Ala Ile Ala Leu Ala
 65 70 75 80
 50 Ala Phe Ser Phe Asn Val Gly His Asp Ala Asn His Asn Ala Tyr Ser
 85 90 95
 Ser Asn Pro His Ile Asn Arg Val Leu Gly Met Thr Tyr Asp Phe Val
 100 105 110
 55 Gly Leu Ser Ser Phe Leu Trp Arg Tyr Arg His Asn Tyr Leu His His
 115 120 125
 Thr Tyr Thr Asn Ile Leu Gly His Asp Val Glu Ile His Gly Asp Gly
 130 135 140
 60 Ala Val Arg Met Ser Pro Glu Gln Glu His Val Gly Ile Tyr Arg Phe
 145 150 155 160

Gln Gln Phe Tyr Ile Trp Gly Leu Tyr Leu Phe Ile Pro Phe Tyr Trp
 165 170 175
 5 Phe Leu Tyr Asp Val Tyr Leu Val Leu Asn Lys Gly Lys Tyr His Asp
 180 185 190
 His Lys Ile Pro Pro Phe Gln Pro Leu Glu Leu Ala Ser Leu Leu Gly
 195 200 205
 10 Ile Lys Leu Leu Trp Leu Gly Tyr Val Phe Gly Leu Pro Leu Ala Leu
 210 215 220
 Gly Phe Ser Ile Pro Glu Val Leu Ile Gly Ala Ser Val Thr Tyr Met
 225 230 235 240
 15 Thr Tyr Gly Ile Val Val Cys Thr Ile Phe Met Leu Ala His Val Leu
 245 250 255
 20 Glu Ser Thr Glu Phe Leu Thr Pro Asp Gly Glu Ser Gly Ala Ile Asp
 260 265 270
 Asp Glu Trp Ala Ile Cys Gln Ile Arg Thr Thr Ala Asn Phe Ala Thr
 275 280 285
 25 Asn Asn Pro Phe Trp Asn Trp Phe Cys Gly Gly Leu Asn His Gln Val
 290 295 300
 Thr His His Leu Phe Pro Asn Ile Cys His Ile His Tyr Pro Gln Leu
 305 310 315 320
 Glu Asn Ile Ile Lys Asp Val Cys Gln Glu Phe Gly Val Glu Tyr Lys
 325 330 335
 35 Val Tyr Pro Thr Phe Lys Ala Ala Ile Ala Ser Asn Tyr Arg Trp Leu
 340 345 350
 Glu Ala Met Gly Lys Ala Ser
 355

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 365 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met Thr Ser Thr Thr Ser Lys Val Thr Phe Gly Lys Ser Ile Gly Phe
 1 5 10 15
 Arg Lys Glu Leu Asn Arg Arg Val Asn Ala Tyr Leu Glu Ala Glu Asn
 20 25 30
 60 Ile Ser Pro Arg Asp Asn Pro Pro Met Tyr Leu Lys Thr Ala Ile Ile
 35 40 45

	Leu	Ala	Trp	Val	Val	Ser	Ala	Trp	Thr	Phe	Val	Val	Phe	Gly	Pro	Asp	
	50						55					60					
5	Val	Leu	Trp	Met	Lys	Leu	Leu	Gly	Cys	Ile	Val	Leu	Gly	Phe	Gly	Val	
	65					70					75					80	
	Ser	Ala	Val	Gly	Phe	Asn	Ile	Ser	His	Asp	Gly	Asn	His	Gly	Gly	Tyr	
					85					90					95		
10	Ser	Lys	Tyr	Gln	Trp	Val	Asn	Tyr	Leu	Ser	Gly	Leu	Thr	His	Asp	Ala	
				100					105					110			
	Ile	Gly	Val	Ser	Ser	Tyr	Leu	Trp	Lys	Phe	Arg	His	Asn	Val	Leu	His	
15			115					120					125				
	His	Thr	Tyr	Thr	Asn	Ile	Leu	Gly	His	Asp	Val	Glu	Ile	His	Gly	Asp	
		130					135					140					
20	Glu	Leu	Val	Arg	Met	Ser	Pro	Ser	Met	Glu	Tyr	Arg	Trp	Tyr	His	Arg	
	145					150					155					160	
	Tyr	Gln	His	Trp	Phe	Ile	Trp	Phe	Val	Tyr	Pro	Phe	Ile	Pro	Tyr	Tyr	
				165						170					175		
25	Trp	Ser	Ile	Ala	Asp	Val	Gln	Thr	Met	Leu	Phe	Lys	Arg	Gln	Tyr	His	
				180					185					190			
	Asp	His	Glu	Ile	Pro	Ser	Pro	Thr	Trp	Val	Asp	Ile	Ala	Thr	Leu	Leu	
30			195					200					205				
	Ala	Phe	Lys	Ala	Phe	Gly	Val	Ala	Val	Phe	Leu	Ile	Ile	Pro	Ile	Ala	
		210					215					220					
35	Val	Gly	Tyr	Ser	Pro	Leu	Glu	Ala	Val	Ile	Gly	Ala	Ser	Ile	Val	Tyr	
	225					230					235					240	
	Met	Thr	His	Gly	Leu	Val	Ala	Cys	Val	Val	Phe	Met	Leu	Ala	His	Val	
				245						250					255		
40	Ile	Glu	Pro	Ala	Glu	Phe	Leu	Asp	Pro	Asp	Asn	Leu	His	Ile	Asp	Asp	
			260						265					270			
	Glu	Trp	Ala	Ile	Ala	Gln	Val	Lys	Thr	Thr	Val	Asp	Phe	Ala	Pro	Asn	
45			275					280					285				
	Asn	Thr	Ile	Ile	Asn	Trp	Tyr	Val	Gly	Gly	Leu	Asn	Tyr	Gln	Thr	Val	
		290					295					300					
50	His	His	Leu	Phe	Pro	His	Ile	Cys	His	Ile	His	Tyr	Pro	Lys	Ile	Ala	
	305					310					315					320	
	Pro	Ile	Leu	Ala	Glu	Val	Cys	Glu	Glu	Phe	Gly	Val	Asn	Tyr	Ala	Val	
				325						330					335		
55	His	Gln	Thr	Phe	Phe	Gly	Ala	Leu	Ala	Ala	Asn	Tyr	Ser	Trp	Leu	Lys	
			340					345						350			
	Lys	Met	Ser	Ile	Asn	Pro	Glu	Thr	Lys	Ala	Ile	Glu	Gln				
60			355					360				365					

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "Synthetic oligonucleotide"

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 21
 (D) OTHER INFORMATION: /number= 1
/note= "N=Inosine or Cytosine"

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 27
 (D) OTHER INFORMATION: /number= 2
/note= "N=Inosine or Cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:
CUACUACUAC UACAYCAYAC NTAYACNAAY AT 32

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid
 (A) DESCRIPTION: /desc = "Synthetic oligonucleotide"

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 13
 (D) OTHER INFORMATION: /number= 1
/note= "N=Inosine or Cytosine"

(ix) FEATURE:
 (A) NAME/KEY: misc_feature
 (B) LOCATION: 19
 (D) OTHER INFORMATION: /number= 2
/note= "N=Inosine or Cytosine"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:
CAUCAUCAUC AUNGGRAANA RRTGRTG 27

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 35 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CCAAGCTTCT GCAGGAGCTC TTTTTTTTTT TTTT 35

10 (2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

(B) TYPE: amino acid

15 (C) STRANDEDNESS: not relevant

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

His Xaa Xaa His His

1 5

25 (2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 5 amino acids

30 (B) TYPE: amino acid

(C) STRANDEDNESS: not relevant

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

35 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Gln Xaa Xaa His His

40 1 5

45 (2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 746 nucleic acids

(B) TYPE: nucleic acid

50 (C) STRANDEDNESS: not relevant

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

55 CGTATGTCAC TCCATTCCAA ACTCGTTCAT GGTATCATAA ATATCAACAC ATTTACGCTC 60

CACTCCTCTA TGGTATTTAC ACACTCAAAT ATCGTACTCA AGATTGGGAA GCTTTTGTA 120

AGGATGGTAA AAATGGTGCA ATTCGTGTTA GTGTCGCCAC AAATTTTCGAT AAGGCCGCTT 180

ACGTCATTGG TAAATTGTCT TTTGTTTTCT TCCGTTTCAT CCTTCCACTC CGTTATCATA 240

60 GCTTACAGA TTTAATTTGT TATTTCCTCA TTGCTGAATT CGTCTTTGGT TGGTATCTCA 300

CAATTAATTT CCAAGTTAGT CATGTCGCTG AAGATCTCAA ATTCTTTGCT ACCCCTGAAA 360

GACCAGATGA ACCATCTCAA ATCAATGAAG ATTGGGCAAT CCTTCAACTT AAAACTACTC 420

AAGATTATGG TCATGGTTCA CTCCTTTGTA CCTTTTITAG TGGTTCCTTA AATCATCAAG 480
 TTGTTTCATCA TTTATTCCCA TCAATTGCTC AAGATTCTTA CCCACAACCTT GTACCAATTG 540
 TAAAAGAAGT TTGTAAAGAA CATAACATTA CTTACCACAT TAAACCAAAC TTCCTGAAG 600
 CTATTATGTC ACACATTAAT TACCTTTACA AAATGGGTAA TGATCCAGAT TATGTTAAAA 660
 5 AACCATTAGC CTCAAAAGAT GATTAAATGA AATAACTTAA AAACCAATTA TTTACTTTTG 720
 ACAAACAGTA ATATTAATAA ATACAA 746

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 227 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Tyr Val Thr Pro Phe Gln Thr Arg Ser Trp Tyr His Lys Tyr Gln
 1 5 10 15
 His Ile Tyr Ala Pro Leu Leu Tyr Gly Ile Tyr Thr Leu Lys Tyr
 20 25 30
 25 Arg Thr Gln Asp Trp Glu Ala Phe Val Lys Asp Gly Lys Asn Gly
 35 40 45
 Ala Ile Arg Val Ser Val Ala Thr Asn Phe Asp Lys Ala Ala Tyr
 50 55 60
 Val Ile Gly Lys Leu Ser Phe Val Phe Phe Arg Phe Ile Leu Pro
 30 65 70 75
 Leu Arg Tyr His Ser Phe Thr Asp Leu Ile Cys Tyr Phe Leu Ile
 80 85 90
 Ala Glu Phe Val Phe Gly Trp Tyr Leu Thr Ile Asn Phe Gln Val
 95 100 105
 35 Ser His Val Ala Glu Asp Leu Lys Phe Phe Ala Thr Pro Glu Arg
 110 115 120
 Pro Asp Glu Pro Ser Gln Ile Asn Glu Asp Trp Ala Ile Leu Gln
 125 130 135
 40 Leu Lys Thr Thr Gln Asp Tyr Gly His Gly Ser Leu Leu Cys Thr
 140 145 150
 Phe Phe Ser Gly Ser Leu Asn His Gln Val Val His His Leu Phe
 155 160 165
 Pro Ser Ile Ala Gln Asp Phe Tyr Pro Gln Leu Val Pro Ile Val
 170 175 180
 45 Lys Glu Val Cys Lys Glu His Asn Ile Thr Tyr His Ile Lys Pro
 185 190 195
 Asn Phe Thr Glu Ala Ile Met Ser His Ile Asn Tyr Leu Tyr Lys
 200 205 210
 50 Met Gly Asn Asp Pro Asp Tyr Val Lys Lys Pro Leu Ala Ser Lys
 215 220 225
 Asp Asp ***

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 494 nucleic acids
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

5 TTTTGAAGG NTCCAAGTTN ACCACGGANT NGGCAAGTTN ACGGGGCGGA AANCGGTTTT 60
 CCCCCCAAGC CTTTTGTCGA CTGGTTCGT GGTGGCTTCC AGTACCAAGT CGACCACCAC 120
 TTATTCCTCA GCCTGCCCCG ACACAATCTG GCCAAGACAC ACGCACTGGT CGAATCGTTC 180
 TGCAAGGAGT GGGGTGTCCA GTACCACGAA GCCGACCTCG TGGACGGGAC CATGGAAGTC 240
 TTGCACCATT TGGGCAGCGT GGCCGCGGAA TTCGTCGTGG ATTTTGTACG CGACGGACCC 300
 10 GCCATGTAAT CGTCGTTCTG GACGATGCAA GGGTTCACGC ACATCTACAC AACTCACTC 360
 ACACAAC TAGTAACTCGT ATAGAATTCG GTGTCGACCT GGACCTTGTT TGA CTGTTG 420
 GGGATAGGGT AGGTAGCGCG ACGCGTGGGT CGNCCCCGGG AATTCTGTGA CCGGTACCTG 480
 GCCCGCTNA AAGT 494

15

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:
 20 (A) LENGTH: 87 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

30 Phe Trp Lys Xxx Pro Ser Xxx Pro Arg Xxx Xxx Gln Val Xxx Gly
 1 5 10 15
 Ala Glu Xxx Gly Phe Pro Pro Lys Pro Phe Val Asp Trp Phe Cys
 20 25 30
 Gly Gly Phe Gln Tyr Gln Val Asp His His Leu Phe Pro Ser Leu
 35 35 40 45
 Pro Arg His Asn Leu Ala Lys Thr His Ala Leu Val Glu Ser Phe
 50 55 60
 Cys Lys Glu Trp Gly Val Gln Tyr His Glu Ala Asp Leu Val Asp
 65 70 75
 Gly Thr Met Glu Val Leu His His Leu Gly Ser Val Ala Gly Glu
 40 65 70 75
 Phe Val Val Asp Phe Val Arg Asp Gly Pro Ala Met
 80 85

45

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:
 50 (A) LENGTH: 520 nucleic acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

55 (ii) MOLECULE TYPE: nucleic acid

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

60 GGATGGAGTT CGTCTGGATC GCTGTGCGCT ACGCGACGTG GTTTAAGCGT CATGGGTGCG 60
 CTTGGGTACA CGCCGGGGCA GTCGTTGGGC ATGTACTTGT GCGCCTTTGG TCTCGGCTGC 120
 ATTTACATT TTCTGCAGTT CGCCGTAAGT CACACCCATT TGCCCGTGAG CAACCCGGAG 180
 GATCAGTGC ATTGGCTCGA GTACGCGCG ACCACACTGT GAACATCAGC ACCAAGTCGT 240

5 GGT TTG TGC AC ATG GTG GATG TCG AACCTCA ACT TTTCAGAT CGAGCACCAC CTTTTCCCA 300
 CGGCGCCCCA GTTCCGTTTC AAGGAGATCA GCCCGCGCGT CGAGGCCCTC TTCAAGCGCC 360
 ACGGTCTCCC TTACTACGAC ATGCCCTACA CGAGCGCCGT CTCCACCACC TTTGCCAACC 420
 TCTACTCCGT CGGCCATTCC GTCGGCGACG CCAAGCGCGA CTAGCCTCTT TTCCTAGACC 480
 TTAATTCCCC ACCCCACCCC ATGTTCTGTC TTCCTCCGC 520

(2) INFORMATION FOR SEQ ID NO:18:

10 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 153 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

20 Met Glu Phe Val Trp Ile Ala Val Arg Tyr Ala Thr Trp Phe Lys
 1 5 10 15
 Arg His Gly Cys Ala Trp Val His Ala Gly Ala Val Val Gly His
 20 25 30
 25 Val Leu Val Arg Leu Trp Ser Arg Leu His Leu His Phe Ser Ala
 35 40 45
 Val Arg Arg Lys Ser His Pro Phe Ala Arg Glu Gln Pro Gly Gly
 50 55 60
 30 Ser Ala Ala Leu Ala Arg Val Arg Ala Asp His Thr Val Asn Ile
 65 70 75
 Ser Thr Lys Ser Trp Phe Val Thr Trp Trp Met Ser Asn Leu Asn
 80 85 90
 Phe Gln Ile Glu His His Leu Phe Pro Thr Ala Pro Gln Phe Arg
 95 100 105
 35 Phe Lys Glu Ile Ser Pro Arg Val Glu Ala Leu Phe Lys Arg His
 110 115 120
 Gly Leu Pro Tyr Tyr Asp Met Pro Tyr Thr Ser Ala Val Ser Thr
 125 130 135
 40 Thr Phe Ala Asn Leu Tyr Ser Val Gly His Ser Val Gly Asp Ala
 140 145 150
 Lys Arg Asp

45 (2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 429 nucleic acids
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: nucleic acid

55 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

60 ACGCGTCCGC CCACGCGTCC GCCGCGAGCA ACTCATCAAG GAAGGCTACT TTGACCCCTC 60
 GCTCCCGCAC ATGACGTACC GCGTGGTCA GATTGTTGTT CTCTTCGTGC TTTCCTTTTG 120
 GCTGATGGGT CAGTCTTCAC CCCTCGCGCT CGCTCTCGGC ATGTCGTCA GCGGCATCTC 180
 TCAGGGTCGC TCGGCTGGG TAATGCATGA GATGGGCCAT GGGTCGTTCA CTGGTGTCTC 240
 TTGGCTTGAC GACCGTTGT GCGAGTTCTT TTACGGCGTT GGTGTGGCA TGAGCGGTCA 300

TTACTGGAAA AACCAGCACA GCAAACACCA CGCAGCGCCA AACCGGCTCG AGCACGATGT 360
 AGATCTCAAC ACCTTGCCAT TGGTGGCCTT CAACGAGCGC GTCGTGCGCA AGGTCCGACC 420

5

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

10

- (A) LENGTH: 125 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

20 Arg Val Arg Pro Arg Val Arg Arg Glu Gln Leu Ile Lys Glu Gly
 1 5 10 15
 Tyr Phe Asp Pro Ser Leu Pro His Met Thr Tyr Arg Val Val Glu
 20 25 30
 Ile Val Val Leu Phe Val Leu Ser Phe Trp Leu Met Gly Gln Ser
 35 40 45
 25 Ser Pro Leu Ala Leu Ala Leu Gly Ile Val Val Ser Gly Ile Ser
 50 55 60
 Gln Gly Arg Cys Gly Trp Val Met His Glu Met Gly His Gly Ser
 65 70 75
 30 Phe Thr Gly Val Ile Trp Leu Asp Asp Arg Leu Cys Glu Phe Phe
 65 70 75
 Tyr Gly Val Gly Cys Gly Met Ser Gly His Tyr Trp Lys Asn Gln
 80 85 90
 His Ser Lys His His Ala Ala Pro Asn Arg Leu Glu His Asp Val
 95 100 105
 35 Asp Leu Asn Thr Leu Pro Leu Val Ala Phe Asn Glu Arg Val Val
 110 115 120
 Arg Lys Val Arg Pro
 125

40

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

45

- (A) LENGTH: 1219 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid (Edited Contig 2692004)

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

55 GCACGCCGAC CGGCGCCGGG AGATCCTGGC AAAGTATCCA GAGATAAAGT CCTTGATGAA 60
 ACCTGATCCC AATTTGATAT GGATTATAAT TATGATGGTT CTCACCCAGT TGGGTGCATT 120
 TTACATAGTA AAAGACTTGG ACTGGAAATG GGTCAATATT GGGGCCTATG CGTTTGGCAG 180
 60 TTGCATTAAC CACTCAATGA CTCTGGCTAT TCATGAGATT GCCCACAATG CTGCCTTTGG 240
 CAACTGCAAA GCAATGTGGA ATCGCTGGTT TGGAATGTTT GCTAATCTTC CTATTGGGAT 300

5 TCCATATTCA ATTCCTTTA AGAGGTATCA CATGGATCAT CATCGGTACC TTGGAGCTGA 360
 TGGCGTCGAT GTAGATATTC CTACCGATTT TGAGGGCTGG TTCTTCTGTA CCGCTTTCAG 420
 AAAGTTTATA TGGGTTATTC TTCAGCCTCT CTTTTATGCC TTTGACCTC TGTCATCAA 480
 CCCCAAACCA ATTACGTATC TGAAGTTAT CAATACCGTG GCACAGGTCA CTTTGTACAT 540
 10 TTTAATTTAT TACTTTTTGG GAATTAAATC CTTAGTCTAC ATGTTGGCAG CATCTTTACT 600
 TGGCCTGGGT TTGCACCCAA TTTCTGGACA TTTTATAGCT GAGCATTACA TGTCTTAAA 660
 GGGTCATGAA ACTTACTCAT ATTATGGGCC TCTGAATTTA CTTACCTTCA ATGTGGGTTA 720
 15 TCATAATGAA CATCATGATT TCCCCAACAT TCCTGGAAAA AGTCTCCAC TGGTGAGGAA 780
 AATAGCAGCT GAATACTATG ACAACCTCCC TCACTACAAT TCCTGGATAA AAGTACTGTA 840
 TGATTTTGTG ATGGATGATA CAATAAGTCC CTA CTCAAGA ATGAAGAGGC ACCAAAAAGG 900
 20 AGAGATGGTG CTGGAGTAAA TATCATTAGT GCCAAAGGGA TTCTTCTCCA AAACCTTAGA 960
 TGATAAAATG GAATTTTGC ATTATTAAAC TTGAGACCAG TGATGCTCAG AAGCTCCCT 1020
 25 GGCACAATTT CAGAGTAAGA GCTCGGTGAT ACCAAGAAGT GAATCTGGCT TTTAAACAGT 1080
 CAGCCTGACT CTGTACTGCT CAGTTTCACT CACAGGAAAC TTGTGACTTG TGTATTATCG 1140
 TCATTGAGGA TGTTTCACTC ATGTCTGTCA TTTTATAAGC ATATCATTTA AAAAGCTTCT 1200
 30 AAAAAGCTAT TTCGCCAGG 1219

(2) INFORMATION FOR SEQ ID NO:22:

35

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 655 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: other nucleic acid (Edited Contig 2153526)

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TTACCTTCTA CGTCCGCTTC TTCCTCACTT ATGTGCCACT ATTGGGGCTG AAAGCTTCCT 60
 50 GGGCCTTTTC TTCATAGTCA GGTTCCTGGA AAGCAACTGG TTTGTGTGGG TGACACAGAT 120
 GAACCATATT CCCATGCACA TTGATCATGA CCGGAACATG GACTGGGTTT CCACCCAGCT 180
 55 CCAGGCCACA TGCAATGTCC ACAAGTCTGC CTTCAATGAC TGGTTCAGTG GACACCTCAA 240
 CTTCCAGATT GAGCACCATC TTTTCCAC GATGCCTCGA CACAATTACC ACAAAGTGGC 300
 TCCCCTGGTG CAGTCCTTGT GTGCCAAGCA TGGCATAGAG TACCAGTCCA AGCCCCTGCT 360
 60 GTCAGCCTTC GCCGACATCA TCCACTCACT AAAGGAGTCA GGGCAGCTCT GGCTAGATGC 420
 CTATCTTCAC CAATAACAAC AGCCACCCTG CCCAGTCTGG AAGAAGAGGA GGAAGACTCT 480
 65 GGAGCCAAGG CAGAGGGGAG CTTGAGGGAC AATGCCACTA TAGTTTAATA CTCAGAGGGG 540

GTTGGGTTTG GGGACATAAA GCCTCTGACT CAAACTCCTC CCTTTTATCT TCTAGCCACA 600
 GTTCTAAGAC CCAAAGTGGG GGGTGGACAC AGAAGTCCCT AGGAGGGAAG GAGCT 655

5

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 304 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

10

(ii) MOLECULE TYPE: other nucleic acid (Edited Contig 3506132)

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GTCTTTTACT TTGGCAATGG CTGGATTCCT ACCCTCATCA CGGCCTTTGT CCTTGCTACC 60
 TCTCAGGCCC AAGCTGGATG GCTGCAACAT GATTATGGCC ACCTGTCTGT CTACAGAAAA 120
 CCCAAGTGGG ACCACCTTGT CCACAAATTC GTCATTGGCC ACTTAAAGGG TGCCTCTGCC 180
 AACTGGTGGG ATCATCGCCA CTTCCAGCAC CACGCCAAGC CTAACATCTT CCACAAGGAT 240
 CCCGATGTGA ACATGCTGCA CGTGTGTTGT CTGGGCGAAT GGCAGCCCAT CGAGTACGGC 300
 AAGA 304

30

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 918 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

35

(ii) MOLECULE TYPE: other nucleic acid (Edited Contig 3854933)

40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CAGGGACCTA CCCCGCGCTA CTTACCTGG GACGAGGTGG CCCAGCGCTC AGGGTGCAGAG 60
 GAGCGGTGGC TAGTGATCGA CCGTAAGGTG TACAACATCA GCGAGTTCAC CCGCCGGCAT 120
 CCAGGGGGCT CCCGGGTCAT CAGCCACTAC GCCGGGCAGG ATGCCACGGA TCCCTTTGTG 180
 GCCTTCCACA TCAACAAGGG CCTTGTGAAG AAGTATATGA ACTCTCTCCT GATTGGAGAA 240
 CTGTCTCCAG AGCAGCCCAG CTTGAGCCC ACCAAGAATA AAGAGCTGAC AGATGAGTTC 300
 CGGGAGCTGC GGGCCACAGT GGAGCGGATG GGGCTCATGA AGGCCAACCA TGTCTTCTTC 360
 CTGCTGTACC TGCTGCACAT CTTGCTGCTG GATGGTGCAG CCTGGCTCAC CCTTTGGGTC 420
 TTTGGGACGT CCTTTTGGC CTTCTCTCTC TGTGCGGTGC TGCTCAGTGC AGTTCAGGCC 480
 CAGGCTGGCT GGCTGCAGCA TGACTTTGGG CACCTGTGCG TCTTCAGCAC CTCAAAGTGG 540
 AACCATCTGC TACATCATTT TGTGATTGGC CACCTGAAGG GGGCCCCCGC CAGTTGGTGG 600
 AACCACATGC ACTTCCAGCA CCATGCCAAG CCCAACTGCT TCCGCAAAGA CCCAGACATC 660

60

AACATGCATC CCTTCTTCTT TGCCTTGGGG AAGATCCTCT CTGTGGAGCT TGGGAAACAG 720
 AAGAAAAAAT ATATGCCGTA CAACCACCAG CACARATACT TCTTCCTAAT TGGGCCCCCA 780
 5 GCCTTGCTGC CTCTCTACTT CCAGTGGTAT ATTTTCTATT TTGTTATCCA GCGAAAGAAG 840
 TGGGTGGACT TGGCCTGGAT CAGCAAACAG GAATACGATG AAGCCGGGCT TCCATTGTCC 900
 10 ACCGCAAATG CTTCTAAA 918

(2) INFORMATION FOR SEQ ID NO:25:

15 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1686 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: other nucleic acid (Edited Contig 2511785)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

25 GCCACTTAAA GGGTGCCTCT GCCAACTGGT GGAATCATCG CCACTTCCAG CACCACGCCA 60
 AGCCTAACAT CTTCCACAAG GATCCCGATG TGAACATGCT GCACGTGTTT GTTCTGGGCG 120
 30 AATGGCAGCC CATCGAGTAC GGCAAGAAGA AGCTGAAATA CCTGCCCTAC AATCACCAGC 180
 ACGAATACTT CTTCTGATT GGGCCGCCGC TGCTCATCCC CATGTATTTT CAGTACCAGA 240
 TCATCATGAC CATGATCGTC CATAAGAACT GGGTGGACCT GGCCTGGGCC GTCAGCTACT 300
 35 ACATCCGGTT CTTTCATCACC TACATCCCTT TCTACGGCAT CCTGGGAGCC CTCCTTTTCC 360
 TCAACTTCAT CAGGTTCTTG GAGAGCCACT GGTGTGTGTG GGTACACAG ATGAATCACA 420
 40 TCGTCATGGA GATTGACCAG GAGGCTTACC GTGACTGGTT CAGTAGCCAG CTGACAGCCA 480
 CCTGCAACGT GGAGCAGTCC TTCTTCAACG ACTGGTTTCTG TGGACACCTT AACTTCCAGA 540
 TTGAGCACCA CCTCTTCCCC ACCATGCCCC GGCACAACTT ACACAAGATC GCCCCGCTGG 600
 45 TGAAGTCTCT ATGTGCCAAG CATGGCATTG AATACCAGGA GAAGCCGCTA CTGAGGGCCC 660
 TGCTGGACAT CATCAGGTCC CTGAAGAAGT CTGGGAAGCT GTGGCTGGAC GCCTACCTTC 720
 50 ACAAATGAAG CCACAGCCCC CGGGACACCG TGGGGAAGGG GTGCAGGTGG GGTGATGGCC 780
 AGAGGAATGA TGGGCTTTTG TTCTGAGGGG TGTCCGAGAG GCTGGTGTAT GCACTGCTCA 840
 CGGACCCCAT GTTGATCTT TCTCCCTTTC TCCTCTCCTT TTTCTCTTCA CATCTCCCCC 900
 55 ATAGCACCCCT GCCCTCATGG GACCTGCCCT CCCTCAGCCG TCAGCCATCA GCCATGGCCC 960
 TCCAGTGCC TCCTAGCCCC TTCTTCCAAG GAGCAGAGAG GTGGCCACCG GGGGTGGCTC 1020
 60 TGTCCTACCT CCACTCTCTG CCCCTAAAGA TGGGAGGAGA CCAGCGGTCC ATGGGTCTGG 1080
 CCTGTGAGTC TCCCCTTGCA GCCTGGTCAC TAGGCATCAC CCCCCTTTG GTTCTTCAGA 1140
 TGCTCTTGGG GTTCATAGGG GCAGGTCCTA GTCGGGCAGG GCCCCTGACC CTCCCGGCTC 1200
 65 GGCTTCACTC TCCCTGACGG CTGCCATTGG TCCACCCTTT CATAGAGAGG CCTGCTTTGT 1260

5 TACAAAGCTC GGGTCTCCCT CCTGCAGCTC GGTAAAGTAC CCGAGGCCTC TCTTAAGATG 1320
 TCCAGGGCCC CAGGCCCCGCG GGCACAGCCA GCCCAAACCT TGGGCCCTGG AAGAGTCCTC 1380
 CACCCCATCA CTAGAGTGCT CTGACCCTGG GCTTTCACGG GCCCATTC ACCGCCTCCC 1440
 CAACTTGAGC CTGTGACCTT GGGACCAAAG GGGGAGTCCC TCGTCTCTTG TGA CTCAGCA 1500
 10 GAGGCAGTGG CCACGTT CAG GGAGGGGCCG GCTGGCCTGG AGGCTCAGCC CACCCTCCAG 1560
 CTTTTCCTCA GGGTGTCTG AGGTCCAAGA TTCTGGAGCA ATCTGACCCT TCTCCAAAGG 1620
 15 CTCTGTTATC AGCTGGGCAG TGCCAGCCAA TCCCTGGCCA TTTGGCCCCA GGGGACGTGG 1680
 GCCCTG 1686

20 (2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1843 base pairs
 (B) TYPE: nucleic acid
 25 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid (Contig 2535)

30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GTCTTTTACT TTGGCAATGG CTGGATTCTT ACCCTCATCA CGGCCTTTGT CCTTGCTACC 60
 35 TCTCAGGCCC AAGCTGGATG GCTGCAACAT GATTATGGCC ACCTGTCTGT CTACAGAAAA 120
 CCCAAGTGGA ACCACCTTGT CCACAAATTC GTCATTGGCC ACTTAAAGGG TGCTCTGCC 180
 AACTGGTGGA ATCATCGCCA CTTCCAGCAC CACGCCAAGC CTAACATCTT CCACAAGGAT 240
 40 CCCGATGTGA ACATGCTGCA CGTGTCTTGT CTGGGCGAAT GGCAGCCCAT CGAGTACGGC 300
 AAGAAGAAGC TGAAATACCT GCCCTACAAT CACCAGCACG AATACTTCTT CCTGATTGGG 360
 45 CCGCCGCTGC TCATCCCCAT GTATTTCCAG TACCAGATCA TCATGACCAT GATCGTCCAT 420
 AAGAACTGGG TGGACCTGGC CTGGGCCGTC AGCTACTACA TCCGGTTCTT CATCACCTAC 480
 ATCCCTTTCT ACGGCATCCT GGGAGCCCTC CTTTTCCTCA ACTTCATCAG GTTCCTGGAG 540
 50 AGCCACTGGT TTGTGTGGGT CACACAGATG AATCACATCG TCATGGAGAT TGACCAGGAG 600
 GCCTACCGTG ACTGGTTCAG TAGCCAGCTG ACAGCCACCT GCAACGTGGA GCAGTCCTTC 660
 55 TTCAACGACT GGTT CAGTGG ACACCTTAAC TTCCAGATTG AGCACCACCT CTCCCCACC 720
 ATGCCCCGGC ACAACTTACA CAAGATCGCC CCGCTGGTGA AGTCTCTATG TGCCAAGCAT 780
 GGCATTGAAT ACCAGGAGAA GCCGCTACTG AGGGCCCTGC TGGACATCAT CAGGTCCCTG 840
 60 AAGAAGTCTG GGAAGCTGTG GCTGGACGCC TACCTTCACA AATGAAGCCA CAGCCCCCGG 900
 GACACCGTGG GGAAGGGGTG CAGGTGGGGT GATGGCCAGA GGAATGATGG GCTTTTGTTC 960
 65 TGAGGGGTGT CCGAGAGGCT GGTGTATGCA CTGCTCACGG ACCCATGTT GGATCTTTCT 1020

CCCTTTCTCC TCTCCTTTTT CTCTTCACAT CTCCCCATA GCACCCTGCC CTCATGGGAC 1080
 CTGCCCTCCC TCAGCCGTCA GCCATCAGCC ATGGCCCTCC CAGTGCCTCC TAGCCCCTTC 1140
 5 TTCCAAGGAG CAGAGAGGTG GCCACCGGGG GTGGCTCTGT CCTACCTCCA CTCTCTGCCC 1200
 CTAAAGATGG GAGGAGACCA GCGGTCCATG GGTCTGGCCT GTGAGTCTCC CCTTGCAGCC 1260
 10 TGGTCACTAG GCATCACCCC CGCTTTGGTT CTTCAGATGC TCTTGGGGTT CATAGGGGCA 1320
 GGTCCTAGTC GGGCAGGGCC CCTGACCCTC CCGGCCTGGC TTCACTCTCC CTGACGGCTG 1380
 CCATTGGTCC ACCCTTTCAT AGAGAGGCCT GCTTTGTTAC AAAGCTCGGG TCTCCCTCCT 1440
 15 GCAGCTCGGT TAAGTACCCG AGGCCTCTCT TAAGATGTCC AGGGCCCCAG GCGCGGGC 1500
 ACAGCCAGCC CAAACCTTGG GCCCTGGAAG AGTCCTCCAC CCCATCACTA GAGTGCTCTG 1560
 20 ACCCTGGGCT TTCACGGGCC CCATTCCACC GCCTCCCCAA CTTGAGCCTG TGACCTTGGG 1620
 ACCAAAGGGG GAGTCCCTCG TCTCTGTGA CTCAGCAGAG GCAGTGGCCA CGTTCAGGGA 1680
 GGGGCCGGCT GGCCTGGAGG CTCAGCCCAC CCTCCAGCTT TTCCTCAGGG TGTCTGAGG 1740
 25 TCCAAGATTC TGGAGCAATC TGACCCTTCT CCAAAGGCTC TGTATCAGC TGGGCAGTGC 1800
 CAGCCAATCC CTGGCCATTT GGCCCCAGGG GACGTGGGCC CTG 1843

30 (2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 2257 base pairs
 35 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid (Edited Contig 253538a)

40 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

CAGGGACCTA CCCCGCGCTA CTTACCTGG GACGAGGTGG CCCAGCGCTC AGGGTGCAG 60
 GAGCGGTGGC TAGTGATCGA CCGTAAGGTG TACAACATCA GCGAGTTCAC CCGCCGGCAT 120
 45 CCAGGGGGCT CCCGGGTCAT CAGCCACTAC GCCGGGCAGG ATGCCACGGA TCCCTTTGTG 180
 GCCTTCCACA TCAACAAGGG CTTGTGAAG AAGTATATGA ACTCTCTCCT GATTGGAGAA 240
 50 CTGTCTCCAG AGCAGCCCAG CTTTGAGCCC ACCAAGAATA AAGAGCTGAC AGATGAGTTC 300
 CGGGAGCTGC GGGCCACAGT GGAGCGGATG GGGCTCATGA AGGCCAACCA TGTCTTCTTC 360
 CTGTGTACC TGCTGCACAT CTTGCTGCTG GATGGTGCAG CCTGGCTCAC CCTTTGGGTC 420
 55 TTTGGGACGT CCTTTTGGC CTTCTCTC TGTGCGGTGC TGCTCAGTGC AGTTCAGCAG 480
 GCCCAAGCTG GATGGCTGCA ACATGATTAT GGCCACCTGT CTGTCTACAG AAAACCCAAG 540
 60 TGGAACCACC TTGTCCACAA ATTCGTCATT GGCCACTTAA AGGGTGCCTC TGCCAAGTGG 600
 TGGAATCATC GCCACTTCCA GCACCACGCC AAGCCTAACA TCTTCCACAA GGATCCCGAT 660
 65 GTGAACATGC TGCACGTGTT TGTCTGGGC GAATGGCAGC CCATCGAGTA CGGCAAGAAG 720

```

AAGCTGAAAT ACCTGCCCTA CAATCACCAG CACGAATACT TCTTCCTGAT TGGGCCGCCG 780
CTGCTCATCC CCATGTATTT CCAGTACCAG ATCATCATGA CCATGATCGT CCATAAGAAC 840
5 TGGGTGGACC TGGCCTGGGC CGTCAGCTAC TACATCCGGT TCTTCATCAC CTACATCCCT 900
TTCTACGGCA TCCTGGGAGC CCTCCTTTTC CTCAACTTCA TCAGGTTCCCT GGAGAGCCAC 960
10 TGGTTTGTGT GGGTCACACA GATGAATCAC ATCGTCATGG AGATTGACCA GGAGGCCTAC 1020
CGTGACTGGT TCAGTAGCCA GCTGACAGCC ACCTGCAACG TGGAGCAGTC CTTCTTCAAC 1080
GACTGGTTCA GTGGACACCT TAACTTCCAG ATTGAGCACC ACCTCTTCCC CACCATGCCC 1140
15 CGGCACAAC TACACAAGAT CGCCCCGTG GTGAAGTCTC TATGTGCCAA GCATGGCATT 1200
GAATACCAGG AGAAGCCGCT ACTGAGGGCC CTGCTGGACA TCATCAGGTC CCTGAAGAAG 1260
20 TCTGGGAAGC TGTGGCTGGA CGCCTACCTT CACAAATGAA GCCACAGCCC CCGGGACACC 1320
GTGGGGAAGG GGTGCAGGTG GGGTGATGGC CAGAGGAATG ATGGGCTTTT GTTCTGAGGG 1380
GTGTCCGAGA GGCTGGTGTA TGCCTGCTC ACGGACCCCA TGTGGATCT TTCTCCCTTT 1440
25 CTCCTCTCCT TTTTCTCTT ACATCTCCCC CATAGCACCC TGCCCTCATG GGACCTGCCC 1500
TCCCTCAGCC GTCAGCCATC AGCCATGGCC CTCCAGTGC CTCCTAGCCC CTTCTTCCAA 1560
30 GGAGCAGAGA GGTGGCCACC GGGGGTGGCT CTGTCTTACC TCCACTCTCT GCCCCTAAAG 1620
ATGGGAGGAG ACCAGCGGTC CATGGGTCTG GCCTGTGAGT CTCCCCTTGC AGCCTGGTCA 1680
CTAGGCATCA CCCCCGCTTT GGTTCCTCAG ATGCTCTTGG GGTTCATAGG GGCAGTCTCT 1740
35 AGTCGGGCAG GGGCCCTGAC CCTCCCGGCC TGGCTTCACT CTCCCTGACG GCTGCCATTG 1800
GTCCACCCTT TCATAGAGAG GCCTGCTTTG TTACAAAGCT CGGGTCTCCC TCCTGCAGCT 1860
40 CGGTTAAGTA CCCGAGGCCT CTCTTAAGAT GTCCAGGGCC CCAGGCCCGC GGGCACAGCC 1920
AGCCCAAACC TTGGGCCCTG GAAGAGTCCT CCACCCATC ACTAGAGTGC TGTGACCCTG 1980
GGCTTTCACG GGCCCCATTC CACCGCCTCC CCAACTTGAG CCTGTGACCT TGGGACCAAA 2040
45 GGGGGAGTCC CTCGTCTCTT GTGACTCAGC AGAGGCAGTG GCCACGTTCA GGGAGGGGCC 2100
GGCTGGCCTG GAGGCTCAGC CCACCCTCCA GCTTTTCCTC AGGGTGTCTT GAGGTCCAAG 2160
ATTCTGGAGC AATCTGACCC TTCTCAAAG GCTCTGTTAT CAGCTGGGCA GTGCCAGCCA 2220
50 ATCCCTGGCC ATTTGGCCCC AGGGGACGTG GGCCCTG 2257

```

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 411 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: amino acid (Translation of Contig 2692004)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

	His	Ala	Asp	Arg	Arg	Arg	Glu	Ile	Leu	Ala	Lys	Tyr	Pro	Glu	Ile	
	1				5					10					15	
5	Lys	Ser	Leu	Met	Lys	Pro	Asp	Pro	Asn	Leu	Ile	Trp	Ile	Ile	Ile	
					20					25					30	
	Met	Met	Val	Leu	Thr	Gln	Leu	Gly	Ala	Phe	Tyr	Ile	Val	Lys	Asp	
					35					40					45	
	Leu	Asp	Trp	Lys	Trp	Val	Ile	Phe	Gly	Ala	Tyr	Ala	Phe	Gly	Ser	
					50					55					60	
10	Cys	Ile	Asn	His	Ser	Met	Thr	Leu	Ala	Ile	His	Glu	Ile	Ala	His	
					65					70					75	
	Asn	Ala	Ala	Phe	Gly	Asn	Cys	Lys	Ala	Met	Trp	Asn	Arg	Trp	Phe	
					80					85					90	
15	Gly	Met	Phe	Ala	Asn	Leu	Pro	Ile	Gly	Ile	Pro	Tyr	Ser	Ile	Ser	
					95					100					105	
	Phe	Lys	Arg	Tyr	His	Met	Asp	His	His	Arg	Tyr	Leu	Gly	Ala	Asp	
					110					115					120	
	Gly	Val	Asp	Val	Asp	Ile	Pro	Thr	Asp	Phe	Glu	Gly	Trp	Phe	Phe	
					125					130					135	
20	Cys	Thr	Ala	Phe	Arg	Lys	Phe	Ile	Trp	Val	Ile	Leu	Gln	Pro	Leu	
					140					145					150	
	Phe	Tyr	Ala	Phe	Arg	Pro	Leu	Phe	Ile	Asn	Pro	Lys	Pro	Ile	Thr	
					155					160					165	
25	Tyr	Leu	Glu	Val	Ile	Asn	Thr	Val	Ala	Gln	Val	Thr	Phe	Asp	Ile	
					170					175					180	
	Leu	Ile	Tyr	Tyr	Phe	Leu	Gly	Ile	Lys	Ser	Leu	Val	Tyr	Met	Leu	
					185					190					195	
	Ala	Ala	Ser	Leu	Leu	Gly	Leu	Gly	Leu	His	Pro	Ile	Ser	Gly	His	
					200					205					210	
30	Phe	Ile	Ala	Glu	His	Tyr	Met	Phe	Leu	Lys	Gly	His	Glu	Thr	Tyr	
					215					220					225	
	Ser	Tyr	Tyr	Gly	Pro	Leu	Asn	Leu	Leu	Thr	Phe	Asn	Val	Gly	Tyr	
					230					235					240	
35	His	Asn	Glu	His	His	Asp	Phe	Pro	Asn	Ile	Pro	Gly	Lys	Ser	Leu	
					245					250					255	
	Pro	Leu	Val	Arg	Lys	Ile	Ala	Ala	Glu	Tyr	Tyr	Asp	Asn	Leu	Pro	
					260					265					270	
	His	Tyr	Asn	Ser	Trp	Ile	Lys	Val	Leu	Tyr	Asp	Phe	Val	Met	Asp	
					275					280					285	
40	Asp	Thr	Ile	Ser	Pro	Tyr	Ser	Arg	Met	Lys	Arg	His	Gln	Lys	Gly	
					290					295					300	
	Glu	Met	Val	Leu	Glu	***	Ile	Ser	Leu	Val	Pro	Lys	Gly	Phe	Phe	
					305					310					315	
45	Ser	Lys	Thr	Leu	Asp	Lys	Met	Glu	Phe	Leu	His	Tyr	***	Thr		
					320					325					330	
	***	Asp	Gln	***	Cys	Ser	Glu	Ala	Pro	Leu	Ala	Gln	Phe	Gln	Ser	
					335					340					345	
	Lys	Ser	Ser	Val	Ile	Pro	Arg	Ser	Glu	Ser	Gly	Phe	***	Thr	Val	
					350					355					360	
50	Ser	Leu	Thr	Leu	Tyr	Cys	Ser	Val	Ser	Leu	Thr	Gly	Asn	Leu	***	
					365					370					375	
	Leu	Val	Tyr	Tyr	Arg	His	***	Gly	Cys	Phe	Thr	His	Val	Cys	His	
					380					385					390	
55	Phe	Ile	Ser	Ile	Ser	Phe	Lys	Lys	Leu	Leu	Lys	Ser	Tyr	Phe	Ala	
					400					405					410	
	Arg															

(2) INFORMATION FOR SEQ ID NO:29:

60

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 218 amino acids
 (B) TYPE: amino acid

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: amino acid (Translation of Contig 2153526)

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

10	Tyr	Leu	Leu	Arg	Pro	Leu	Leu	Pro	His	Leu	Cys	Ala	Thr	Ile	Gly	1	5	10	15
	Ala	Glu	Ser	Phe	Leu	Gly	Leu	Phe	Phe	Ile	Val	Arg	Phe	Leu	Glu	20	25	30	
	Ser	Asn	Trp	Phe	Val	Trp	Val	Thr	Gln	Met	Asn	His	Ile	Pro	Met	35	40	45	
15	His	Ile	Asp	His	Asp	Arg	Asn	Met	Asp	Trp	Val	Ser	Thr	Gln	Leu	50	55	60	
	Gln	Ala	Thr	Cys	Asn	Val	His	Lys	Ser	Ala	Phe	Asn	Asp	Trp	Phe	65	70	75	
	Ser	Gly	His	Leu	Asn	Phe	Gln	Ile	Glu	His	His	Leu	Phe	Pro	Thr	80	85	90	
20	Met	Pro	Arg	His	Asn	Tyr	His	Lys	Val	Ala	Pro	Leu	Val	Gln	Ser	95	100	105	
	Leu	Cys	Ala	Lys	His	Gly	Ile	Glu	Tyr	Gln	Ser	Lys	Pro	Leu	Leu	110	115	120	
25	Ser	Ala	Phe	Ala	Asp	Ile	Ile	His	Ser	Leu	Lys	Glu	Ser	Gly	Gln	125	130	135	
	Leu	Trp	Leu	Asp	Ala	Tyr	Leu	His	Gln	***	Gln	Gln	Pro	Pro	Cys	140	145	150	
	Pro	Val	Trp	Lys	Lys	Arg	Arg	Lys	Thr	Leu	Glu	Pro	Arg	Gln	Arg	155	160	165	
30	Gly	Ala	***	Gly	Thr	Met	Pro	Leu	***	Phe	Asn	Thr	Gln	Arg	Gly	170	175	180	
	Leu	Gly	Leu	Gly	Thr	***	Ser	Leu	***	Leu	Lys	Leu	Leu	Pro	Phe	185	190	195	
35	Ile	Phe	***	Pro	Gln	Phe	***	Asp	Pro	Lys	Trp	Gly	Val	Asp	Thr	200	205	210	
	Glu	Val	Pro	Arg	Arg	Glu	Gly	Ala								215			

40

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 71 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

45

50

(ii) MOLECULE TYPE: amino acid (Translation of Contig 3506132)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

55

60

	Val	Phe	Tyr	Phe	Gly	Asn	Gly	Trp	Ile	Pro	Thr	Leu	Ile	Thr	Ala	1	5	10	15
	Phe	Val	Leu	Ala	Thr	Ser	Gln	Ala	Gln	Ala	Gly	Trp	Leu	Gln	His	20	25	30	
	Asp	Tyr	Gly	His	Leu	Ser	Val	Tyr	Arg	Lys	Pro	Lys	Trp	Asn	His	35	40	45	
	Leu	Val	His	Lys	Phe	Val	Ile	Gly	His	Leu	Lys	Gly	Ala	Ser	Ala				

305

5 (2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 566 amino acids

(B) TYPE: amino acid

10 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: amino acid (Translation of Contig 2511785)

15 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

	His	Leu	Lys	Gly	Ala	Ser	Ala	Asn	Trp	Trp	Asn	His	Arg	His	Phe
	1				5					10					15
20	Gln	His	His	Ala	Lys	Pro	Asn	Ile	Phe	His	Lys	Asp	Pro	Asp	Val
					20					25					30
	Asn	Met	Leu	His	Val	Phe	Val	Leu	Gly	Glu	Trp	Gln	Pro	Ile	Glu
					35					40					45
	Tyr	Gly	Lys	Lys	Lys	Leu	Lys	Tyr	Leu	Pro	Tyr	Asn	His	Gln	His
					50					55					60
25	Glu	Tyr	Phe	Phe	Leu	Ile	Gly	Pro	Pro	Leu	Leu	Ile	Pro	Met	Tyr
					65					70					75
	Phe	Gln	Tyr	Gln	Ile	Ile	Met	Thr	Met	Ile	Val	His	Lys	Asn	Trp
					80					85					90
30	Val	Asp	Leu	Ala	Trp	Ala	Val	Ser	Tyr	Tyr	Ile	Arg	Phe	Phe	Ile
					95					100					105
	Thr	Tyr	Ile	Pro	Phe	Tyr	Gly	Ile	Leu	Gly	Ala	Leu	Leu	Phe	Leu
					110					115					120
	Asn	Phe	Ile	Arg	Phe	Leu	Glu	Ser	His	Trp	Phe	Val	Trp	Val	Thr
					125					130					135
35	Gln	Met	Asn	His	Ile	Val	Met	Glu	Ile	Asp	Gln	Glu	Ala	Tyr	Arg
					140					145					150
	Asp	Trp	Phe	Ser	Ser	Gln	Leu	Thr	Ala	Thr	Cys	Asn	Val	Glu	Gln
					155					160					165
40	Ser	Phe	Phe	Asn	Asp	Trp	Phe	Ser	Gly	His	Leu	Asn	Phe	Gln	Ile
					170					175					180
	Glu	His	His	Leu	Phe	Pro	Thr	Met	Pro	Arg	His	Asn	Leu	His	Lys
					185					190					195
	Ile	Ala	Pro	Leu	Val	Lys	Ser	Leu	Cys	Ala	Lys	His	Gly	Ile	Glu
					200					205					210
45	Tyr	Gln	Glu	Lys	Pro	Leu	Leu	Arg	Ala	Leu	Leu	Asp	Ile	Ile	Arg
					215					220					225
	Ser	Leu	Lys	Lys	Ser	Gly	Lys	Leu	Trp	Leu	Asp	Ala	Tyr	Leu	His
					230					235					240
	Lys	***	Ser	His	Ser	Pro	Arg	Asp	Thr	Val	Gly	Lys	Gly	Cys	Arg
50					245					250					255
	Trp	Gly	Asp	Gly	Gln	Arg	Asn	Asp	Gly	Leu	Leu	Phe	***	Gly	Val
					260					265					270
	Ser	Glu	Arg	Leu	Val	Tyr	Ala	Leu	Leu	Thr	Asp	Pro	Met	Leu	Asp
					275					280					285
55	Leu	Ser	Pro	Phe	Leu	Leu	Ser	Phe	Phe	Ser	Ser	His	Leu	Pro	His
					290					295					300
	Ser	Thr	Leu	Pro	Ser	Trp	Asp	Leu	Pro	Ser	Leu	Ser	Arg	Gln	Pro
					305					310					315
	Ser	Ala	Met	Ala	Leu	Pro	Val	Pro	Pro	Ser	Pro	Phe	Phe	Gln	Gly
60					320					325					330
	Ala	Glu	Arg	Trp	Pro	Pro	Gly	Val	Ala	Leu	Ser	Tyr	Leu	His	Ser
					335					340					345

5 Leu Pro Leu Lys Met Gly Gly Asp Gln Arg Ser Met Gly Leu Ala
 350 355 360
 Cys Glu Ser Pro Leu Ala Ala Trp Ser Leu Gly Ile Thr Pro Ala
 365 370 375
 10 Leu Val Leu Gln Met Leu Leu Gly Phe Ile Gly Ala Gly Pro Ser
 380 385 390
 Arg Ala Gly Pro Leu Thr Leu Pro Ala Trp Leu His Ser Pro ***
 400 405 410
 Arg Leu Pro Leu Val His Pro Phe Ile Glu Arg Pro Ala Leu Leu
 415 420 425
 Gln Ser Ser Gly Leu Pro Pro Ala Ala Arg Leu Ser Thr Arg Gly
 430 435 440
 Leu Ser *** Asp Val Gln Gly Pro Arg Pro Ala Gly Thr Ala Ser
 445 450 455
 15 Pro Asn Leu Gly Pro Trp Lys Ser Pro Pro Pro His His *** Ser
 460 465 470
 Ala Leu Thr Leu Gly Phe His Gly Pro His Ser Thr Ala Ser Pro
 475 480 485
 Thr *** Ala Cys Asp Leu Gly Thr Lys Gly Gly Val Pro Arg Leu
 490 495 500
 20 Leu *** Leu Ser Arg Gly Ser Gly His Val Gln Gly Gly Ala Gly
 505 510 515
 Trp Pro Gly Gly Ser Ala His Pro Pro Ala Phe Pro Gln Gly Val
 520 525 530
 25 Leu Arg Ser Lys Ile Leu Glu Gln Ser Asp Pro Ser Pro Lys Ala
 535 540 545
 Leu Leu Ser Ala Gly Gln Cys Gln Pro Ile Pro Gly His Leu Ala
 550 555 560
 30 Pro Gly Asp Val Gly Pro Xxx
 565

(2) INFORMATION FOR SEQ ID NO:33:

35 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 619 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 40 (ii) MOLECULE TYPE: amino acid (Translation of Contig 2535)
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

45 Val Phe Tyr Phe Gly Asn Gly Trp Ile Pro Thr Leu Ile Thr Ala
 1 5 10 15
 50 Phe Val Leu Ala Thr Ser Gln Ala Gln Ala Gly Trp Leu Gln His
 20 25 30
 Asp Tyr Gly His Leu Ser Val Tyr Arg Lys Pro Lys Trp Asn His
 35 40 45
 Leu Val His Lys Phe Val Ile Gly His Leu Lys Gly Ala Ser Ala
 50 55 60
 55 Asn Trp Trp Asn His Arg His Phe Gln His His Ala Lys Pro Asn
 65 70 75
 Ile Phe His Lys Asp Pro Asp Val Asn Met Leu His Val Phe Val
 80 85 90
 60 Leu Gly Glu Trp Gln Pro Ile Glu Tyr Gly Lys Lys Lys Leu Lys
 95 100 105
 Tyr Leu Pro Tyr Asn His Gln His Glu Tyr Phe Phe Leu Ile Gly
 110 115 120

	Pro	Pro	Leu	Leu	Ile	Pro	Met	Tyr	Phe	Gln	Tyr	Gln	Ile	Ile	Met
					125					130					135
	Thr	Met	Ile	Val	His	Lys	Asn	Trp	Val	Asp	Leu	Ala	Trp	Ala	Val
					140					145					150
5	Ser	Tyr	Tyr	Ile	Arg	Phe	Phe	Ile	Thr	Tyr	Ile	Pro	Phe	Tyr	Gly
					155					160					165
	Ile	Leu	Gly	Ala	Leu	Leu	Phe	Leu	Asn	Phe	Ile	Arg	Phe	Leu	Glu
					170					175					180
10	Ser	His	Trp	Phe	Val	Trp	Val	Thr	Gln	Met	Asn	His	Ile	Val	Met
					185					190					195
	Glu	Ile	Asp	Gln	Glu	Ala	Tyr	Arg	Asp	Trp	Phe	Ser	Ser	Gln	Leu
					200					205					210
	Thr	Ala	Thr	Cys	Asn	Val	Glu	Gln	Ser	Phe	Phe	Asn	Asp	Trp	Phe
					215					220					225
15	Ser	Gly	His	Leu	Asn	Phe	Gln	Ile	Glu	His	His	Leu	Phe	Pro	Thr
					230					235					240
	Met	Pro	Arg	His	Asn	Leu	His	Lys	Ile	Ala	Pro	Leu	Val	Lys	Ser
					245					250					255
20	Leu	Cys	Ala	Lys	His	Gly	Ile	Glu	Tyr	Gln	Glu	Lys	Pro	Leu	Leu
					260					265					270
	Arg	Ala	Leu	Leu	Asp	Ile	Ile	Arg	Ser	Leu	Lys	Lys	Ser	Gly	Lys
					275					280					285
	Leu	Trp	Leu	Asp	Ala	Tyr	Leu	His	Lys	***	Ser	His	Ser	Pro	Arg
					290					295					300
25	Asp	Thr	Val	Gly	Lys	Gly	Cys	Arg	Trp	Gly	Asp	Gly	Gln	Arg	Asn
					305					310					315
	Asp	Gly	Leu	Leu	Phe	***	Gly	Val	Ser	Glu	Arg	Leu	Val	Tyr	Ala
					320					325					330
30	Leu	Leu	Thr	Asp	Pro	Met	Leu	Asp	Leu	Ser	Pro	Phe	Leu	Leu	Ser
					335					340					345
	Phe	Phe	Ser	Ser	His	Leu	Pro	His	Ser	Thr	Leu	Pro	Ser	Trp	Asp
					350					355					360
	Leu	Pro	Ser	Leu	Ser	Arg	Gln	Pro	Ser	Ala	Met	Ala	Leu	Pro	Val
					365					370					375
35	Pro	Pro	Ser	Pro	Phe	Phe	Gln	Gly	Ala	Glu	Arg	Trp	Pro	Pro	Gly
					380					385					390
	Val	Ala	Leu	Ser	Tyr	Leu	His	Ser	Leu	Pro	Leu	Lys	Met	Gly	Gly
					400					405					410
40	Asp	Gln	Arg	Ser	Met	Gly	Leu	Ala	Cys	Glu	Ser	Pro	Leu	Ala	Ala
					415					420					425
	Trp	Ser	Leu	Gly	Ile	Thr	Pro	Ala	Leu	Val	Leu	Gln	Met	Leu	Leu
					430					435					440
	Gly	Phe	Ile	Gly	Ala	Gly	Pro	Ser	Arg	Ala	Gly	Pro	Leu	Thr	Leu
					445					450					455
45	Pro	Ala	Trp	Leu	His	Ser	Pro	***	Arg	Leu	Pro	Leu	Val	His	Pro
					460					465					470
	Phe	Ile	Glu	Arg	Pro	Ala	Leu	Leu	Gln	Ser	Ser	Gly	Leu	Pro	Pro
					475					480					485
50	Ala	Ala	Arg	Leu	Ser	Thr	Arg	Gly	Leu	Ser	***	Asp	Val	Gln	Gly
					490					495					500
	Pro	Arg	Pro	Ala	Gly	Thr	Ala	Ser	Pro	Asn	Leu	Gly	Pro	Trp	Lys
					505					510					515
	Ser	Pro	Pro	Pro	His	His	***	Ser	Ala	Leu	Thr	Leu	Gly	Phe	His
					520					525					530
55	Gly	Pro	His	Ser	Thr	Ala	Ser	Pro	Thr	***	Ala	Cys	Asp	Leu	Gly
					535					540					545
	Thr	Lys	Gly	Gly	Val	Pro	Arg	Leu	Leu	***	Leu	Ser	Arg	Gly	Ser
					550					555					560
60	Gly	His	Val	Gln	Gly	Gly	Ala	Gly	Trp	Pro	Gly	Gly	Ser	Ala	His
					565					570					575
	Pro	Pro	Ala	Phe	Pro	Gln	Gly	Val	Leu	Arg	Ser	Lys	Ile	Leu	Glu
					580					585					590

Gln	Ser	Asp	Pro	Ser	Pro	Lys	Ala	Leu	Leu	Ser	Ala	Gly	Gln	Cys
				595					600					605
Gln	Pro	Ile	Pro	Gly	His	Leu	Ala	Pro	Gly	Asp	Val	Gly	Pro	Xxx
				610					615					620

5

(2) INFORMATION FOR SEQ ID NO:34:

10

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 757 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

15

(ii) MOLECULE TYPE: amino acid (Translation of Contig 253538a)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

20

Gln	Gly	Pro	Thr	Pro	Arg	Tyr	Phe	Thr	Trp	Asp	Glu	Val	Ala	Gln
1				5					10					15
Arg	Ser	Gly	Cys	Glu	Glu	Arg	Trp	Leu	Val	Ile	Asp	Arg	Lys	Val
				20					25					30
25	Tyr	Asn	Ile	Ser	Glu	Phe	Thr	Arg	Arg	His	Pro	Gly	Gly	Ser
				35					40					45
Val	Ile	Ser	His	Tyr	Ala	Gly	Gln	Asp	Ala	Thr	Asp	Pro	Phe	Val
				50					55					60
Ala	Phe	His	Ile	Asn	Lys	Gly	Leu	Val	Lys	Lys	Tyr	Met	Asn	Ser
				65					70					75
30	Leu	Leu	Ile	Gly	Glu	Leu	Ser	Pro	Glu	Gln	Pro	Ser	Phe	Glu
				80					85					90
Thr	Lys	Asn	Lys	Glu	Leu	Thr	Asp	Glu	Phe	Arg	Glu	Leu	Arg	Ala
				95					100					105
35	Thr	Val	Glu	Arg	Met	Gly	Leu	Met	Lys	Ala	Asn	His	Val	Phe
				110					115					120
Leu	Leu	Tyr	Leu	Leu	His	Ile	Leu	Leu	Leu	Asp	Gly	Ala	Ala	Trp
				125					130					135
Leu	Thr	Leu	Trp	Val	Phe	Gly	Thr	Ser	Phe	Leu	Pro	Phe	Leu	Leu
				140					145					150
40	Cys	Ala	Val	Leu	Leu	Ser	Ala	Val	Gln	Gln	Ala	Gln	Ala	Gly
				155					160					165
Leu	Gln	His	Asp	Tyr	Gly	His	Leu	Ser	Val	Tyr	Arg	Lys	Pro	Lys
				170					175					180
45	Trp	Asn	His	Leu	Val	His	Lys	Phe	Val	Ile	Gly	His	Leu	Lys
				185					190					195
Ala	Ser	Ala	Asn	Trp	Trp	Asn	His	Arg	His	Phe	Gln	His	His	Ala
				200					205					210
Lys	Pro	Asn	Ile	Phe	His	Lys	Asp	Pro	Asp	Val	Asn	Met	Leu	His
				215					220					225
50	Val	Phe	Val	Leu	Gly	Glu	Trp	Gln	Pro	Ile	Glu	Tyr	Gly	Lys
				230					235					240
Lys	Leu	Lys	Tyr	Leu	Pro	Tyr	Asn	His	Gln	His	Glu	Tyr	Phe	Phe
				245					250					255
55	Leu	Ile	Gly	Pro	Pro	Leu	Leu	Ile	Pro	Met	Tyr	Phe	Gln	Tyr
				260					265					270
Ile	Ile	Met	Thr	Met	Ile	Val	His	Lys	Asn	Trp	Val	Asp	Leu	Ala
				275					280					285
Trp	Ala	Val	Ser	Tyr	Tyr	Ile	Arg	Phe	Phe	Ile	Thr	Tyr	Ile	Pro
				290					295					300
60	Phe	Tyr	Gly	Ile	Leu	Gly	Ala	Leu	Leu	Phe	Leu	Asn	Phe	Ile
				305					310					315

	Phe	Leu	Glu	Ser	His	Trp	Phe	Val	Trp	Val	Thr	Gln	Met	Asn	His	
					320					325					330	
	Ile	Val	Met	Glu	Ile	Asp	Gln	Glu	Ala	Tyr	Arg	Asp	Trp	Phe	Ser	
					335					340					345	
5	Ser	Gln	Leu	Thr	Ala	Thr	Cys	Asn	Val	Glu	Gln	Ser	Phe	Phe	Asn	
					350					355					360	
	Asp	Trp	Phe	Ser	Gly	His	Leu	Asn	Phe	Gln	Ile	Glu	His	His	Leu	
					365					370					375	
10	Phe	Pro	Thr	Met	Pro	Arg	His	Asn	Leu	His	Lys	Ile	Ala	Pro	Leu	
					380					385					390	
	Val	Lys	Ser	Leu	Cys	Ala	Lys	His	Gly	Ile	Glu	Tyr	Gln	Glu	Lys	
					400					405					410	
	Pro	Leu	Leu	Arg	Ala	Leu	Leu	Asp	Ile	Ile	Arg	Ser	Leu	Lys	Lys	
					415					420					425	
15	Ser	Gly	Lys	Leu	Trp	Leu	Asp	Ala	Tyr	Leu	His	Lys	***	Ser	His	
					430					435					440	
	Ser	Pro	Arg	Asp	Thr	Val	Gly	Lys	Gly	Cys	Arg	Trp	Gly	Asp	Gly	
					445					450					455	
20	Gln	Arg	Asn	Asp	Gly	Leu	Leu	Phe	***	Gly	Val	Ser	Glu	Arg	Leu	
					460					465					470	
	Val	Tyr	Ala	Leu	Leu	Thr	Asp	Pro	Met	Leu	Asp	Leu	Ser	Pro	Phe	
					475					480					485	
	Leu	Leu	Ser	Phe	Phe	Ser	Ser	His	Leu	Pro	His	Ser	Thr	Leu	Pro	
					490					495					500	
25	Ser	Trp	Asp	Leu	Pro	Ser	Leu	Ser	Arg	Gln	Pro	Ser	Ala	Met	Ala	
					505					510					515	
	Leu	Pro	Val	Pro	Pro	Ser	Pro	Phe	Phe	Gln	Gly	Ala	Glu	Arg	Trp	
					520					525					530	
30	Pro	Pro	Gly	Val	Ala	Leu	Ser	Tyr	Leu	His	Ser	Leu	Pro	Leu	Lys	
					535					540					545	
	Met	Gly	Gly	Asp	Gln	Arg	Ser	Met	Gly	Leu	Ala	Cys	Glu	Ser	Pro	
					550					555					560	
	Leu	Ala	Ala	Trp	Ser	Leu	Gly	Ile	Thr	Pro	Ala	Leu	Val	Leu	Gln	
					565					570					575	
35	Met	Leu	Leu	Gly	Phe	Ile	Gly	Ala	Gly	Pro	Ser	Arg	Ala	Gly	Pro	
					580					585					590	
	Leu	Thr	Leu	Pro	Ala	Trp	Leu	His	Ser	Pro	***	Arg	Leu	Pro	Leu	
					595					600					605	
40	Val	His	Pro	Phe	Ile	Glu	Arg	Pro	Ala	Leu	Leu	Gln	Ser	Ser	Gly	
					610					615					620	
	Leu	Pro	Pro	Ala	Ala	Arg	Leu	Ser	Thr	Arg	Gly	Leu	Ser	***	Asp	
					625					630					635	
	Val	Gln	Gly	Pro	Arg	Pro	Ala	Gly	Thr	Ala	Ser	Pro	Asn	Leu	Gly	
					640					645					650	
45	Pro	Trp	Lys	Ser	Pro	Pro	Pro	His	His	***	Ser	Ala	Leu	Thr	Leu	
					655					660					665	
	Gly	Phe	His	Gly	Pro	His	Ser	Thr	Ala	Ser	Pro	Thr	***	Ala	Cys	
					670					675					680	
50	Asp	Leu	Gly	Thr	Lys	Gly	Gly	Val	Pro	Arg	Leu	Leu	***	Leu	Ser	
					685					690					695	
	Arg	Gly	Ser	Gly	His	Val	Gln	Gly	Gly	Ala	Gly	Trp	Pro	Gly	Gly	
					700					705					710	
	Ser	Ala	His	Pro	Pro	Ala	Phe	Pro	Gln	Gly	Val	Leu	Arg	Ser	Lys	
					715					720					725	
55	Ile	Leu	Glu	Gln	Ser	Asp	Pro	Ser	Pro	Lys	Ala	Leu	Leu	Ser	Ala	
					730					735					740	
	Gly	Gln	Cys	Gln	Pro	Ile	Pro	Gly	His	Leu	Ala	Pro	Gly	Asp	Val	
					745					750					755	
60	Gly	Pro	Xxx													

What is claimed is:

1. An isolated nucleic acid comprising:
a nucleotide sequence depicted in a SEQ ID NO. 1
2. A polypeptide encoded by said nucleic acid of claim 1.
- 5 3. A purified or isolated polypeptide comprising an amino acid sequence depicted in SEQ ID NO: 2.
4. An isolated nucleic acid encoding the polypeptide of SEQ ID NO: 2.
5. An isolated nucleic acid comprising:
10 a nucleotide sequence which encodes a polypeptide that desaturates a fatty acid molecule at carbon 5 from the carboxyl end of said fatty acid molecule.
6. The isolated nucleic acid according to Claim 5, wherein said nucleotide sequence is derived from eukaryotic cell.
- 15 7. The isolated nucleic acid according to Claim 6, wherein said eukaryotic cell is a fungal cell.
8. The isolated nucleic acid according to Claim 7, wherein said fungal cell is of the genus *Mortierella*.
9. The isolated nucleic acid according to Claim 8, wherein said
20 *Mortierella* cell is of the species *Mortierella alpina*.
10. The isolated nucleic acid according to Claim 5, wherein said nucleotide sequence anneals to a nucleotide sequence depicted in SEQ ID NO: 1.
11. The nucleic acid of claim 10, wherein said nucleotide sequence
25 encodes an amino acid sequence depicted in SEQ ID NO: 2.
12. The nucleic acid of claim 11, wherein said amino acid sequence depicted in SEQ ID NO: 2 is selected from the group consisting of amino acid residues 30-38, 41-44, 171-175, 203-212, and 387-394.

13. An isolated or purified polypeptide which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of said fatty acid molecule.

14. An isolated nucleic acid comprising:

5 a nucleotide sequence which is substantially identical to a sequence of at least 50 nucleotides in SEQ ID NO 1.

15. An isolated nucleic acid sequence having at least about 50% identity to SEQ ID NO 1.

16. A nucleic acid construct comprising:

10 a nucleotide sequence depicted in a SEQ ID NO: 1 linked to a heterologous nucleic acid.

17. A nucleic acid construct comprising:

a nucleotide sequence depicted in a SEQ ID NO: 1 operably linked to a promoter.

15 18. The nucleic acid construct of claim 17, wherein said promoter is functional in a microbial cell.

19. The nucleic acid construct of claim 18, wherein said microbial cell is a yeast cell.

20. The nucleic acid construct of claim 17, wherein said nucleotide sequence is derived from a fungus.

20 21. The nucleic acid according to Claim 19, wherein said fungus is of the genus *Mortierella*.

22. The nucleic acid according to Claim 20, wherein said fungus is of the species *Mortierella alpina*.

23. A nucleic acid construct comprising:

25 a nucleotide sequence which encodes a polypeptide comprising an amino acid sequence which corresponds to or is complementary to an amino acid sequence depicted in SEQ ID NO: 2, wherein said nucleotide sequence is operably linked to a promoter which is functional in a host cell, wherein said nucleotide

sequence encodes a polypeptide which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of a fatty acid molecule.

24. A nucleic acid construct comprising:

5 a nucleotide sequence which encodes a functionally active $\Delta 5$ -desaturase, said desaturase having an amino acid sequence which corresponds to or is complementary to all of or a portion of an amino acid sequence depicted in a SEQ ID NO: 2, wherein said nucleotide sequence is operably linked to a promoter functional in a host cell.

25. A recombinant yeast cell comprising:

10 a nucleic construct according to Claim 23 or Claim 24.

26. The recombinant yeast cell according to Claim 25, wherein said yeast cell is a *Saccharomyces* cell.

27. A host cell comprising:

15 at least one copy of a nucleotide sequence which encodes a polypeptide which converts dihomo- γ -linolenic acid to arachidonic acid, wherein said microbial cell or an ancestor of said microbial cell was transformed with a vector comprising said nucleotide sequence, and wherein said nucleotide sequence is operably linked to a promoter functional in said host cell.

20 28. The microbial cell according to Claim 27, wherein said cell is a host cell selected from the group consisting of a fungal cell and an algal cell.

29. The microbial cell according to Claim 28, wherein said fungal cell is a yeast cell and said algae cell is marine algal cell.

25 30. The microbial cell according to Claim 27, wherein said cell is enriched for 20:3 fatty acids as compared to a host cell which is devoid of said nucleotide sequence.

31. The microbial cell according to Claim 27, wherein said cell is enriched for 20:4 or ω -3 20:4 fatty acids as compared to a host cell which is devoid of said DNA sequence.

32. The microbial cell according to Claim 27, wherein said cell is enriched for 20:5 fatty acids as compared to a host cell which is devoid of said DNA sequence.

5 33. The microbial cell according to Claim 27, wherein said cell has an altered amount of 20:3 (8, 11, 14) fatty acid as compared to an untransformed microbial cell.

34. A method for production of arachidonic acid in a microbial cell culture, said method comprising:

10 growing a microbial cell culture having a plurality of microbial cells, wherein said microbial cells or ancestors of said microbial cells were transformed with a vector comprising one or more nucleic acids having a nucleotide sequence which encodes a polypeptide which converts dihomo- γ -linolenic acid to arachidonic acid, wherein said one or more nucleic acids are operably linked to a promoter, under conditions wherein said one or more
15 nucleic acids are expressed and arachidonic acid is produced in said microbial cell culture.

35. The method of Claim 34, wherein said polypeptide is an enzyme which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of said fatty acid molecule.

20 36. The method of Claim 34, wherein said nucleotide sequence is derived from a *Mortierella species*.

37. The method according to Claim 34, wherein said dihomo- γ -linolenic acid is exogenously supplied.

25 38. The method according to Claim 34, wherein said microbial cells are yeast cells.

39. The method according to Claim 38, wherein said yeast cells are *Saccharomyces species* cells.

40. The method according to Claim 34, wherein said conditions are inducible.

41. A recombinant yeast cell which converts greater than about 5% of a 20:3 fatty acid to a 20:4 fatty acid.

42. A nucleic acid probe comprising:
a nucleotide sequence as represented by SEQ ID NO:1.

5 43. A host cell comprising:
a nucleic acid construct according to Claim 23 or Claim 24.

44. A host cell comprising:
a vector which includes a nucleic acid which encodes a fatty acid
desaturase derived from *Mortierella alpina*, wherein said fatty acid desaturase
10 comprises an amino acid sequence represented by SEQ ID NO:2, wherein said
nucleic acid is operably linked to a promoter.

45. The host cell according to Claim 44, wherein said host cell is a
eukaryotic cell.

46. The host cell according to Claim 45, wherein said eukaryotic cell
15 is selected from the group consisting of a mammalian cell, a plant cell, a fungal
cell, an avian cell and an algal cell.

47. The host cell according to Claim 45, wherein said host cell
contains dihomo-gamma-linolenic acid.

48. The host cell according to Claim 45, wherein said host cell
20 contains EPA.

49. The host cell according to Claim 44, wherein said promoter is
exogenously supplied.

50. A method for desaturating a dihomo- γ -linolenic acid, said
method comprising:
25 culturing a recombinant microbial cell according to Claim 37, under
conditions suitable for expression of polypeptide encoded by said nucleic acid,
wherein said host cell further comprises a fatty acid substrate of said
polypeptide.

51. A fatty acid desaturated by the method according to Claim 50.

52. An oil comprising a fatty acid according to Claim 51.

53. A method for obtaining altered long chain polyunsaturated fatty acid biosynthesis comprising the steps of:

5 growing a microbe having cells which contain a transgene which encodes a transgene expression product which desaturates a fatty acid molecule at carbon 5 from the carboxyl end of said fatty acid molecule, wherein said transgene is operably associated with an expression control sequence, under conditions whereby said transgene is expressed, whereby long chain
10 polyunsaturated fatty acid biosynthesis in said cells is altered.

54. A method for obtaining altered long chain polyunsaturated fatty acid biosynthesis comprising the steps of:

growing a microbe having cells which contain a transgene, derived from a fungus or algae, which encodes a transgene expression product which
15 desaturates a fatty acid molecule at carbon 5 from the carboxyl end of said fatty acid molecule, wherein said transgene is operably associated with an expression control sequence, under conditions whereby said one or more transgenes is expressed, whereby long chain polyunsaturated fatty acid biosynthesis in said cells is altered.

20 55. The method according to claims 53 or 54, wherein said long chain polyunsaturated fatty acid is selected from the group consisting of ARA, DGLA and EPA.

56. A microbial oil or fraction thereof produced according to the method of claims 53 or 54.

25 57. A method of treating or preventing malnutrition comprising administering said microbial oil of claim 56 to a patient in need of said treatment or prevention in an amount sufficient to effect said treatment or prevention.

58. A pharmaceutical composition comprising said microbial oil or fraction of claim 56 and a pharmaceutically acceptable carrier.

59. The pharmaceutical composition of claim 58, wherein said pharmaceutical composition is in the form of a solid or a liquid.

60. The pharmaceutical composition of claim 59, wherein said pharmaceutical composition is in a capsule or tablet form.

5 61. The pharmaceutical composition of claim 58 further comprising at least one nutrient selected from the group consisting of a vitamin, a mineral, a carbohydrate, a sugar, an amino acid, a free fatty acid, a phospholipid, an antioxidant, and a phenolic compound.

62. A nutritional formula comprising said microbial oil or fraction
10 thereof of claim 56.

63. The nutritional formula of claim 62, wherein said nutritional formula is selected from the group consisting of an infant formula, a dietary supplement, and a dietary substitute.

64. The nutritional formula of claim 63, wherein said infant formula,
15 dietary supplement or dietary supplement is in the form of a liquid or a solid.

65. An infant formula comprising said microbial oil or fraction thereof of claim 56.

66. The infant formula of claim 65 further comprising at least one macronutrient selected from the group consisting of coconut oil, soy oil, canola
20 oil, mono- and diglycerides, glucose, edible lactose, electrodialysed whey, electrodialysed skim milk, milk whey, soy protein, and other protein hydrolysates.

67. The infant formula of claim 66 further comprising at least one vitamin selected from the group consisting of Vitamins A, C, D, E, and B
25 complex; and at least one mineral selected from the group consisting of calcium, magnesium, zinc, manganese, sodium, potassium, phosphorus, copper, chloride, iodine, selenium, and iron.

68. A dietary supplement comprising said microbial oil or fraction thereof of claim 56.

69. The dietary supplement of claim 68 further comprising at least one macronutrient selected from the group consisting of coconut oil, soy oil, canola oil, mono- and diglycerides, glucose, edible lactose, electro dialysed whey, electro dialysed skim milk, milk whey, soy protein, and other protein hydrolysates.

70. The dietary supplement of claim 69 further comprising at least one vitamin selected from the group consisting of Vitamins A, C, D, E, and B complex; and at least one mineral selected from the group consisting of calcium, magnesium, zinc, manganese, sodium, potassium, phosphorus, copper, chloride, iodine, selenium, and iron.

71. The dietary supplement of claim 68 or claim 70, wherein said dietary supplement is administered to a human or an animal.

72. A dietary substitute comprising said microbial oil or fraction thereof of claim 56.

73. The dietary substitute of claim 72 further comprising at least one macronutrient selected from the group consisting of coconut oil, soy oil, canola oil, mono- and diglycerides, glucose, edible lactose, electro dialysed whey, electro dialysed skim milk, milk whey, soy protein, and other protein hydrolysates.

74. The dietary substitute of claim 73 further comprising at least one vitamin selected from the group consisting of Vitamins A, C, D, E, and B complex; and at least one mineral selected from the group consisting of calcium, magnesium, zinc, manganese, sodium, potassium, phosphorus, copper, chloride, iodine, selenium, and iron.

75. The dietary substitute of claim 72 or claim 74, wherein said dietary substitute is administered to a human or animal.

76. A method of treating a patient having a condition caused by insufficient intake or production of polyunsaturated fatty acids comprising administering to said patient said dietary substitute of claim 72 or said dietary supplement of claim 68 in an amount sufficient to effect said treatment.

77. The method of claim 72, wherein said dietary substitute or said dietary supplement is administered enterally or parenterally.

78. A cosmetic comprising said microbial oil or fraction thereof of claim 56.

5 79. The cosmetic of claim 78, wherein said cosmetic is applied topically.

80. The pharmaceutical composition of claim 58, wherein said pharmaceutical composition is administered to a human or an animal.

10 81. An animal feed comprising said microbial oil or fraction thereof of claim 56.

82. The method of claim 54 wherein said fungus is *Mortierella species*.

83. The method of claim 82 wherein said fungus is *Mortierella alpina*.

15 84. An isolated nucleotide sequence selected from the group consisting of SEQ ID NO:13 and SEQ ID NO:15.

85. An isolated nucleotide sequence from the group consisting of SEQ ID NO:7 and SEQ ID NO:19.

20 86. An isolated nucleotide sequence comprising a nucleotide sequence selected from the group consisting of: SEQ ID NO:13; SEQ ID NO:15; SEQ ID NO:17; SEQ ID NO:19; SEQ ID NO:21; SEQ ID NO:22; SEQ ID NO:23; SEQ ID NO:24; SEQ ID NO:25; SEQ ID NO:26 and SEQ ID NO:27.

25 87. An isolated peptide sequence comprising a peptide sequence selected from the group consisting of: SEQ ID NO:14; SEQ ID NO:16; SEQ ID NO:18; SEQ ID NO:20; SEQ ID NO:28; SEQ ID NO:29; SEQ ID NO:30; SEQ ID NO:31; SEQ ID NO:32; SEQ ID NO:33 and SEQ ID NO:34.

88. Purified polypeptides produced from the nucleotide sequences of claims 84-86.




FIG. 1

PUFA PATHWAYS




FIG. 2

FIG. 3A

GCTTCCTCCA GTTCATCCTC CATTCGCCA CCTGCATTCT TTACGACCGT TAAGCAAG
 60 *
 ATG GGA ACG GAC CAA GGA AAA ACC TTC ACC TGG GAA GAG CTG GCG GCC
 Met Gly Thr Asp Gln Gly Lys Thr Phe Thr Trp Glu Glu Leu Ala Ala
 120 *
 CAT AAC ACC AAG GAC GAC CTA CTC TTG GCC ATC CGC GGC AGG GTG TAC
 His Asn Thr Lys Asp Asp Leu Leu Leu Ala Ile Arg Gly Arg Val Tyr
 180 *
 GAT GTC ACA AAG TTC TTG AGC CGC CAT CCT GGT GGA GTG GAC ACT CTC
 Asp Val Thr Lys Phe Leu Ser Arg His Pro Gly Gly Val Asp Thr Leu
 240 *
 CTG CTC GGA GCT GGC CGA GAT GTT ACT CCG GTC TTT GAG ATG TAT CAC
 Leu Leu Gly Ala Gly Arg Asp Val Thr Pro Val Phe Glu Met Tyr His
 GCG TTT GGG GCT GCA GAT GCC ATT ATG AAG AAC TAC TAT GTC GGT ACA
 Ala Phe Gly Ala Ala Asp Ala Ile Met Lys Lys Tyr Tyr Val Gly Thr
 300 *
 CTG GTC TCG AAT GAG CTG CCC ATC TTC CCG GAG CCA ACG GTG TTC CAC
 Leu Val Ser Asn Glu Leu Pro Ile Phe Pro Glu Pro Thr Val Phe His
 360 *
 AAA ACC ATC AAG ACG AGA GTC GAG GGC TAC TTT ACG GAT CGG AAC ATT
 Lys Thr Ile Lys Thr Arg Val Glu Gly Tyr Phe Thr Asp Arg Asn Ile

FIG. 3B

```

420 *
GAT CCC AAG AAT AGA CCA GAG ATC TGG GGA CGA TAC GCT CTT ATC TTT
Asp Pro Lys Asn Arg Pro Glu Ile Trp Gly Arg Tyr Ala Leu Ile Phe

480 *
GGA TCC TTG ATC GCT TCC TAC TAC GCG CAG CTC TTT GTG CCT TTC GTT
Gly Ser Leu Ile Ala Ser Tyr Tyr Ala Gln Leu Phe Val Pro Phe Val

540 *
GTC GAA CGC ACA TGG CTT CAG GTG GTG TTT GCA ATC ATC ATG GGA TTT
Val Glu Arg Thr Trp Leu Gln Val Val Phe Ala Ile Ile Met Gly Phe

600 *
GCG TGC GCA CAA GTC GGA CTC AAC CCT CTT CAT GAT GCG TCT CAC TTT
Ala Cys Ala Gln Val Gly Leu Asn Pro Leu His Asp Ala Ser His Phe

660 *
TCA GTG ACC CAC AAC CCC ACT GTC TGG AAG ATT CTG GGA GCC ACG CAC
Ser Val Thr His Asn Pro Thr Val Trp Lys Ile Leu Gly Ala Thr His

720 *
GAC TTT TTC AAC GGA GCA TCG TAC CTG GTG TGG ATG TAC CAA CAT ATG
Asp Phe Phe Asn Gly Ala Ser Tyr Leu Val Trp Met Tyr Gln His Met

CTC GGC CAT CAC CCC TAC ACC AAC ATT GCT GGA GCA GAT CCC GAC GTG
Leu Gly His His Pro Tyr Thr Asn Ile Ala Gly Ala Asp Pro Asp Val

```

FIG. 3C

TCG ACG TCT GAG CCC GAT GTT CGT CGT ATC AAG CCC AAC CAA AAG TGG
 Ser Thr Ser Glu Pro Asp Val Arg Arg Ile Lys Pro Asn Gln Lys Trp
 780 *

TTT GTC AAC CAC ATC AAC CAG CAC ATG TTT GTT CCT TTC CTG TAC GGA
 Phe Val Asn His Ile Asn Gln His Met Phe Val Pro Phe Leu Tyr Gly
 840 *

CTG CTG GCG TTC AAG GTG GCG ATT CAG GAC ATC AAC ATT TTG TAC TTT
 Leu Leu Ala Phe Lys Val Arg Ile Gln Asp Ile Asn Ile Leu Tyr Phe
 900 *

GTC AAG ACC AAT GAC GCT ATT CGT GTC AAT CCC ATC TCG ACA TGG CAC
 Val Lys Thr Asn Asp Ala Ile Arg Val Asn Pro Ile Ser Thr Trp His
 960 *

ACT GTG ATG TTC TGG GCG GGC AAG GCT TTC TTT GTC TGG TAT CGC CTG
 Thr Val Met Phe Trp Gly Gly Lys Ala Phe Phe Val Trp Tyr Arg Leu
 1020 *

ATT GTT CCC CTG CAG TAT CTG CCC CTG GGC AAG GTG CTG CTC TTG TTC
 Ile Val Pro Leu Gln Tyr Leu Pro Leu Gly Lys Val Leu Leu Phe
 1080 *

ACG GTC GCG GAC ATG GTG TCG TCT TAC TGG CTG GCG CTG ACC TTC CAG
 Thr Val Ala Asp Met Val Ser Ser Tyr Trp Leu Ala Leu Thr Phe Gln
 1140 *

6/17

FIG. 3D

1080 *
 GCG AAC CAC GTT GTT GAG GAA GTT CAG TGG CCG TTG CCT GAC GAG AAC
 Ala Asn His Val Val Glu Glu Val Gln Trp Pro Leu Pro Asp Glu Asn

1140 *
 GGG ATC ATC CAA AAG GAC TGG GCA GCT ATG CAG GTC GAG ACT ACG CAG
 Gly Ile Ile Gln Lys Asp Trp Ala Ala Met Gln Val Glu Thr Thr Gln

1200 *
 GAT TAC GCA CAC GAT TCG CAC CTC TGG ACC AGC ATC ACT GGC AGC TTG
 Asp Tyr Ala His Asp Ser His Leu Trp Thr Ser Ile Thr Gly Ser Leu

1260 *
 AAC TAC CAG GCT GTG CAC CAT CTG TTC CCC AAC GTG TCG CAG CAC CAT
 Asn Tyr Gln Ala Val His His Leu Phe Pro Asn Val Ser Gln His His

1320 *
 TAT CCC GAT ATT CTG GCC ATC ATC AAG AAC ACC TGC AGC GAG TAC AAG
 Tyr Pro Asp Ile Leu Ala Ile Ile Lys Asn Thr Cys Ser Glu Tyr Lys

1380 *
 GTT CCA TAC CTT GTC AAG GAT ACG TTT TGG CAA GCA TTT GCT TCA CAT
 Val Pro Tyr Leu Val Lys Asp Thr Phe Trp Gln Ala Phe Ala Ser His

1440 *
 TTG GAG CAC TTG CGT GTT CTT GGA CTC CGT CCC AAG GAA GAG TAGA
 Leu Glu His Leu Arg Val Leu Gly Leu Arg Pro Lys Glu Glu

AGAAAAAAG CGCCGAATGA AGTATTGCC CCTTTTCTC CAAGAATGCC AAAAGGAGAT
 CAAGTGGACA TTCTCTATGA AGA

FIG. 4

10	20	30	40	50	60
LHHTYTNIAG ADPDVSTSEP DVRRIPKNQK WFNHINQHM FVPFLYGLLA FKVRIQDINI					
70	80	90	100	110	120
LYFVKTNDAI RVNPISTWHT VMFWGGKAFF VWYRLIVPLQ YLPLGKVLLL FTVADMVSSY					
130	140	150	160	170	180
WLALTFQANY VVEEVQWPLP DENGIIQKDW AAMQVETTQD YAHDSHLWTS ITGSLNYQXV					
HHLFPH					

8/17

1A29	MGIDQG-KTFTM	70
1A524	MAAPSVRTEFRAEVLNABALN	71
BorD6	MA--AQIKKYITSDKPKGQWIS	72
Sy6803D6	ML-TAE-RI	73
SpID6	MTSTTS-DV	74
1A29	FEHYHAFCAADAAMKXYVVG	80
1A524	FDTHFP-EAAMETLANPXYV	81
BorD6	FAVAP-P-ASTWKNLDKFFTC	82
Sy6803D6	---	83
SpID6	---	84
1A29	SLIASYAAQLFVFPFVVER	150
1A524	IWGL--STVIVAKWGQTS	151
BorD6	FIAMLFAMSVYGVLFCE	152
Sy6803D6	WLFSAW--AFVLFAPVIF	153
SpID6	WVVSAM--TFVVF	154
1A29	YLVWWMXQ-HHLLGHHPT	220
1A524	SSHWKDKHNT-HHAA	221
BorD6	SIGWKKWNHN-AHHI	222
Sy6803D6	FEI-MRRR-HNYLHHT	223
SpID6	YI-MKFR-HNVLHHT	224
1A29	LLAF--KVRIQDINILY	290
1A524	ILSLARLSWCLQSL	291
BorD6	IMCAARLNMYVQSLI	292
Sy6803D6	FIPF--YWFLYV	293
SpID6	FIPY--YWSI	294
1A29	ADNVSSYWLALTFQ	360
1A524	SQAVCGNLLAVQFSL	361
BorD6	SLVGTG-MQVQFSL	362
Sy6803D6	TYHTYGVVCTIFM	363
SpID6	VYVTHGLVACVVE	364
1A29	LFPNVYSQHXYRDI	430
1A524	LFFSHMPRNFNSKI	431
BorD6	LFRKMPRCNLKLS	432
Sy6803D6	LFPNICHIIHY	433
SpID6	LFFHICHIIHY	434

FIG. 5

9/17

Effect of Timing of Substrate Addition on Expression of *Mortierella* Delta-5-desaturase Gene in Yeast (SC334) (Induction Temperature 15 C)




FIG. 6

10/17

Effect of Concentration of Inducer (Galactose) on Expression of *Mortierella* Delta-5-desaturase Gene in Yeast (SC334) (Induction Temperature 15 C)




FIG. 7

11/17

Effect of Induction Temperature on Expression of *Mortierella* Delta-5-desaturase Gene in Yeast (Strain SC334)




FIG. 8

12/17

Effect of Yeast Strain on Expression of *Mortierella* Delta-5-desaturase Gene (Induction Temperature 15 C)




FIG. 9

Effect of Yeast Strain on Expression of *Mortierella* Delta-5-desaturase Gene (Induction Temperature 30 C)

WO 98/46765

PCT/US98/07422

13/17




FIG. 10

Expression of delta-5-desaturase in DBY746 versus SC334




FIG. 11

Effect of Medium Composition and Temperature on Expression of delta-5-desaturase Activity in Two Host Strains (SC334 and INVSC1)




FIG. 12

16/17

FastA Match of ma29 and contig 253538a

SCORES Initl: 117 Initn: 225 Opt: 256
 Smith-Waterman score: 408; 27.0% identity in 441 aa overlap

	10	20	30	40	50
ma29gcg.pep	MGTDQGKT---	FTWEELAAHNTKDDLLLAIRGRVYDVT	KFLSRHPGGVD	TLLLGAGRDVT	
		: : :	: : : :	:	:
253538a	QGTPRYFTWDEVAQRSGCEERWLVIDRKVYNISEFTRRHPGGSRVISHYAGQDAT				
	10	20	30	40	50
	60	70	80	90	100
ma29gcg.pep	PVFEMYHAF-GAADA	IMKKYYVGT	LVSNELPIFPEPTV	FHKTIKTRVEGYFTDRNIDPKN	
	:	: :	:	:	: :
253538a	DPFVAFHINKGLVKKYMN	SLIGEL-SPEQPSF-EPTKNKELTDEFREL	RATVERMGLMK		
	60	70	80	90	100
	120	130	140	150	160
ma29gcg.pep	RPEIWGRYALIFGSLIASYYAQLFVPFV	VERTWLQVVF-AIIMGFACAQVGLNPLHDASH			
	: : :	: : :	: : : :	: :	:
253538a	ANHVF--FLLYLLHILLDGA	AWLTWLVFGTSFLPFLCAVLLSAVQAQAGWLQ-HDYGH			
	120	130	140	150	160
	180	190	200	210	220
ma29gcg.pep	FSVTHNPTVWKILGATHDF----	FNGASYLVWMYQHMLGHH	PHYTNIA	GADPDVSTSE---	
	: :	: :	: :	:	:
253538a	LSVYRKPK-WNHL--VHKFVIGHLKGASAN	WWNHRH-FQHHAKPNI	FHKDPD	VNMLHVFV	
	180	190	200	210	220
	230	240	250	260	270
ma29gcg.pep	----PDVRRIPKNQKWF-VNHINQHMFV--	PFLYGLLAFKVRIQDINILYFVKTNDAIRV			
	: : :	: : :	: :	: :	: :
253538a	LGEWQPIEYGKKKLKYL	PYNHQHEFFLIGPPLIPMYFQYQI----	IMTMIVHKNWVDL		
	230	240	250	260	270
	290	300	310	320	330
ma29gcg.pep	NPISTWHTVMFWGGKAFFVWYRLIV	PLQYLPLGKVLLLFTVADMVSSYWLALT	FQANHV		
	: : :	: : :	: : :	: : :	: :
253538a	----AWAVSYYI--RFFITY--	IPF-YGILG-ALLFLNFIRFLESHWFVWVTQMNHIV			
	290	300	310	320	330
	350	360	370	380	390
ma29gcg.pep	EEVQWPLPDENGIIQKDWAA	MQVETT----	QDYAHD	SHLWTSITGSLNYQAVHHLF	PNVS
	:	: : :	: :	:	:
253538a	MEI-----DQEAY--RDWFSSQLTATCNVEQSFFND--	WFS--GHLNFQIEHHLFPTMP			
	340	350	360	370	
	400	410	420	430	440
ma29gcg.pep	QHHPDILAIKNTCSEYKVPYLVKDTFWQAFASHLEHLRVLGLRPKEEX				
	: :	: : :	:		
253538a	RHNLHKIAPLVKSLCAKHGIEYQEKPLLRALLDI	IRSLKKS	GKLWLDAYLHKX		
	380	390	400	410	420

Figure 13

17/17

FastA Match of ma524 and contig 253538a

SCORES Init1: 231 Initn: 499 Opt: 401
 Smith-Waterman score: 620; 27.3% identity in 455 aa overlap

```

      10      20      30      40      50      59
ma524gcg.pep MAAAPSVRTFTRAEVLNAEALNEGKKDAEAPFLMIIDNKVYDVREFVDPHPGGSVILTH-
      | : | | | | : : : : : : : : | | | : | : | | | : : |
253538a      QGPTPRYFTWDEV-----AQRSGCEERWLVIDRKVYNISEFTRRRHPGGSRVISHY
      10      20      30      40      50

      60      70      80      90      100     110
ma524gcg.pep VGKDGTDVFDITFHEAAW--ETLANFYVGDIIDE---SDRDIKNDDFAAEVRKLRTLFSQL
      : : : : | : | : : : : : : : : : : | : | : : | : | : : : :
253538a      AGQDATDPFVAFHINKGLVKKYMNSLLIGELSPQPSFEPTKKNKELTDEFRELRTATVERM
      60      70      80      90      100     110

      120     130     140     150     160     170
ma524gcg.pep GYYDSSKAYYAFKVSFNLCTIWLSTVIVAKWGQTSTLANVLSAALLGLFWQQCGWLHADF
      | : : : : : : : | : | : : : : | | | : | : | : | | | | :
253538a      GLMKANHVFFLLYLLHILLLDGAANLTLWVFG-TSFLPFLICAVLLSAVQAQAGWLQHDY
      120     130     140     150     160

      180     190     200     210     220     230
ma524gcg.pep LHHQVFQDRFWGDLFGAFLGGVCQGFSSSWKKDKHNTTHHAAPNVHGEDPDIDITHPLLTWS
      | : : : : | : | : | : : | : | : | : | : | : | : | : : |
253538a      GHLSVYRKPKNNHLVHKFVIGHLKGASANWNNHRHFQHHAKPNIFHKDPDVN---ML---
      170     180     190     200     210     220

      240     250     260     270     280     290
ma524gcg.pep EHALEMFSIDVPDEELTRMWSRPMVLNQITWFFYPILS---FARLSWCLQSILFVLPNGQAH
      | : : : : | : : : : | : : : | : : : : | : : : : | : : : : |
253538a      -HVF-VLGEWQPIEYGGKKLKYLPYNHQQEYFFLIGPPLLIPMYQYQIIMIMI---VH
      230     240     250     260     270

      300     310     320     330     340     349
ma524gcg.pep KPSGARVPISLVEQLSLAMHWIWLATMFLFIK--DPVNMLVYFLVSQAVCCNLLAIVFS
      | : : : : | : : : : | : : : : | : : : : | : : : : | : : : : |
253538a      K-----NWDLAWAVSYIRFFITYIPFYGILGALLFLNIFRFLSHWFVWVITQ
      280     290     300     310     320

      350     360     370     380     390     400     409
ma524gcg.pep LNHNGMPVISKEEAVDMDFFTKQIITGRDVHPLGFANWFTGGLNYQIEHHLFSPMPRHNF
      : | | : : : | : | : : : : : : : : : : | : | : | : | : | : | :
253538a      MNHIVMEI---DQEAYR-DWFSSQLTATCNVEQSFFNDWFSGHILNFQIEHHLFPTMPRHNL
      330     340     350     360     370     380

      410     420     430     440     450
ma524gcg.pep SKIQPAVETLCKKYNVRYHTTGMIEGTAEVFSRLNEVSKAASKMGKAQX
      | | | : : | : : : : : : : : : : : : : : : : | : : : :
253538a      HKIAPLVKSLCAKHGIEYQEKPLLRLLDIIRSLKSGKLWLDAYLHKX
      390     400     410     420     430

```

Figure 14

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/07422

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/53 C12N15/82 C12N5/10 C12P7/64 C11B1/00
A61K31/20 A23L1/30 A23K1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C12P C11B A61K A23L A23K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	COVELLO P. ET AL.: "Functional expression of the extraplastidial Arabidopsis thaliana oleate desaturase gene (FAD2) in Saccharomyces cerevisiae" PLANT PHYSIOLOGY, vol. 111, no. 1, May 1996, pages 223-226, XP002075211 see the whole document ---	1-83
A	WO 93 06712 A (RHONE POULENC AGROCHIMIE) 15 April 1993 cited in the application see the whole document ---	1-83
A	WO 94 18337 A (MONSANTO CO ;UNIV MICHIGAN (US); GIBSON SUSAN IRMA (US); KISHORE G) 18 August 1994 * see the whole document * ---	1-83
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 September 1998

Date of mailing of the international search report

21/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Kania, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/07422

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 21022 A (RHONE POULENC AGROCHIMIE) 11 July 1996 cited in the application * see the whole document *	1-83
A	EP 0 561 569 A (LUBRIZOL CORP) 22 September 1993 cited in the application see the whole document	1-83
A	SPYCHALLA J. ET AL.: "Identification of an animal w3 fatty acid desaturase by heterologous expression in Arabidopsis" PNAS, U.S.A., vol. 94, no. 4, 18 February 1997, pages 1142-1147, XP002076628 * see esp. discussion *	1-83
A	HILLIER L. ET AL.: "The WashU-Merck EST Project, AC W49761" EMBL DATABASE, 30 May 1996, XP002076629 Heidelberg * corresponding to SEQ ID NO: 21 * see the whole document	86
A	NATHANS J.: "Adult human retina cDNA, AC W28140" EMBL DATABASE, 14 May 1996, XP002076630 Heidelberg * corresponding to SEQ ID NO: 22 * see the whole document	86
A	HILLIER L. ET AL.: "The WashU-Merck EST Project, AC W67716" EMBL DATABASE, 16 June 1996, XP002076631 Heidelberg * corresponding to SEQ ID NO: 24 * see the whole document	86
A	HILLIER L. ET AL.: "The WashU-Merck EST Project, AC H17219" EMBL DATABASE, 1 July 1995, XP002076632 Heidelberg * corresponding to SEQ ID NO: 25, 27 * see the whole document	86

-/--

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/07422

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HILLIER L. ET AL.: "The WashU-Merck EST Project, AC H19385" EMBL DATABASE, 7 July 1995, XP002076633 Heidelberg * corresponding to SEQ ID NO: 26 * see the whole document ---	86
P,X	YOSHINOR. ET AL.: "AC C25549" EMBL DATABASE, 24 July 1997, XP002076634 Heidelberg * corresponding to SEQ ID NO: 13 * see the whole document ---	84,86-88
P,X	CADENA D. ET AL.: "AC AF002668" EMBL DATABASE, 4 July 1997, XP002076635 Heidelberg * corresponding to SEQ ID NO: 21 * see the whole document ---	86-88
T	MICHAELSON L. ET AL.: "Isolation of a delta5-fatty acid desaturase gene from Mortierella alpina" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, no. 30, 24 July 1998, pages 19055-19059, XP002076636 see the whole document -----	1-83

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 07422

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 57, 76, 77
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/ US 98 /07422

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-83

Nucleic acids, polypeptides, constructs comprising delta-5 desaturase according to SEQ ID NO: 1,2 derived from the fungus *Mortierella alpina*. Recombinant host cells comprising said nucleic acids or constructs.

Methods for the production of arachidonic acid in a microbial cell comprising a cell containing a vector encoding an enzyme activity which converts dihomo- γ -linolenic acid to arachidonic acid, preferentially a delta-5 desaturase, more preferentially from *Mortierella*. A recombinant yeast cell converting more than 5% of a 20:3 fatty acid to a 20:4 fatty acid.

Methods for desaturating dihomo- γ -linolenic acid using said microbial cells, fatty acids and oils obtained thereby.

Methods for obtaining altered long chain polyunsaturated fatty acid biosynthesis using microbes comprising delta-5 desaturase derived from fungi or algae.

Microbial oils derived from thereof and their use for therapeutical, nutritional, and cosmetical purposes, as well as products derived therefrom.

2. Claims: 84,86-88 partially

An isolated sequence comprising the sequence of SEQ ID NO: 13, purified polypeptides produced thereof, esp. comprising SEQ ID NO: 14.

3. Claims: 84, 86-88 partially

An isolated nucleotide comprising the sequence of SEQ ID NO: 15, purified polypeptides produced thereof, esp. comprising SEQ ID NO: 16.

4. Claims: 85 completely, 86-88 partially

An isolated nucleotide sequence consisting of SEQ ID NO: 17 and SEQ ID NO: 19, purified polypeptides produced therefrom, esp. comprising SEQ ID NO: 18 and SEQ ID NO: 20.

./.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/07422

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9306712 A	15-04-1993	AU 667848 B	18-04-1996
		AU 2881292 A	03-05-1993
		BG 98695 A	31-05-1995
		BR 9206613 A	11-04-1995
		CA 2120629 A	15-04-1993
		CN 1072722 A	02-06-1993
		CN 1174236 A	25-02-1998
		CZ 9400817 A	13-09-1995
		EP 0666918 A	16-08-1995
		HU 69781 A	28-09-1995
		JP 7503605 T	20-04-1995
		MX 9205820 A	01-04-1993
		NZ 244685 A	27-06-1994
		US 5552306 A	03-09-1996
		US 5614393 A	25-03-1997
		US 5689050 A	18-11-1997
		US 5663068 A	02-09-1997
		US 5789220 A	04-08-1998
		ZA 9207777 A	21-04-1993
WO 9418337 A	18-08-1994	EP 0684998 A	06-12-1995
		JP 8506490 T	16-07-1996
WO 9621022 A	11-07-1996	US 5614393 A	25-03-1997
		AU 4673596 A	24-07-1996
		CA 2207906 A	11-07-1996
		CN 1177379 A	25-03-1998
		EP 0801680 A	22-10-1997
		US 5789220 A	04-08-1998
EP 0561569 A	22-09-1993	AU 3516793 A	16-09-1993
		CA 2092661 A	14-09-1993
		JP 6014667 A	25-01-1994
		US 5777201 A	07-07-1998